

**WELL-POSEDNESS FOR A CAUCHY
FRACTIONAL DIFFERENTIAL PROBLEM WITH
HILFER TYPE FRACTIONAL DERIVATIVE**

BY

MOHAMMED DAHAN AHMED KASSIM

A Thesis Presented to the
DEANSHIP OF GRADUATE STUDIES

KING FAHD UNIVERSITY OF PETROLEUM & MINERALS

DHAHRAN, SAUDI ARABIA

In Partial Fulfillment of the
Requirements for the Degree of

MASTER OF SCIENCE

In

MATHEMATICS

MAY, 2011


KING FAHD UNIVERSITY OF PETROLEUM & MINERALS

DHAHRAN 31261, SAUDI ARABIA

DEANSHIP OF GRADUATE STUDIES

This thesis, written by **MOHAMMED DAHAN AHMED KASSIM** under the direction of his thesis advisor and approved by his thesis committee, has been presented to and accepted by the Dean of Graduate Studies, in partial fulfillment of the requirements for the degree of **MASTER OF SCIENCE IN MATHEMATICS**.

Thesis Committee




Prof. Nasser-eddine Tatar (Advisor)




Dr. Khaled Furati (Co- Advisor)



Prof. Salim Messaoudi (Member)



Dr. M. Tahir Mustafa (Member)



Dr. Kassem Mustapha (Member)



Dr. Hattan Tawfiq
Department Chairman



Dr. Salam A. Zummo
Dean of Graduate Studies



الإهداء

أهدي هذا العمل

إلى والدي العزيزين ثمرة من ثمار تربيتهن وغرسهم

إلى زوجتي العزيزة و اولادي.... وفاءً لهم ولصبرهم وتضحيتهم

إلى أخواتي رداً لبعض معروفهن وإحسانهن

DEDICATED

TO

MY PARENTS AND FAMILY

ACKNOWLEDGMENT

I am most grateful to Almighty ALLAH, the Beneficent, the Merciful, for enabling me to complete this work. Peace and blessings of ALLAH be upon his Last messenger Mohammed (Sallallah-Alaihe-Wasallam) and his family, who guided us to the right path.

First and the foremost acknowledgments are due to King Fahd University of Petroleum and Minerals and to the Department of Mathematics and Statistics for supporting my research work.

I wish to express my deep appreciation to **Professor Nasser-eddine Tatar** who served as my major advisor, for his guidance and patience through the preparation of thesis. His continuous support and encouragement can never be forgotten. My thanks are extended to my Co-advisor **Dr. Khaled Furati** who has guided me with his dedicated attention, expertise and knowledge throughout this research. I also wish to thank my committee members, **Prof. Salim Messaoudi, Dr. M. Tahir Mustafa** and **Dr. Kassem Mustapha** for their constructive guidance and support.

I would like also to thank the Chairman of the Mathematics Department Dr. Hattan Tawfiq for providing all the available facilities. I am also grateful to all faculty members for their encouragement and their direct or indirect help. Finally, I thank my family for their patience and encouragement during my years of study.

TABLE OF CONTENTS

THESIS ABSTRACT (ENGLISH)	VI
THESIS ABSTRACT (ARABIC)	VII
CHAPTER 1: INTRODUCTION	1
1.1 WHAT IS FRACTIONAL CALCULUS.....	2
1.2 FIRST CONTRIBUTIONS	2
1.3 FIRST BOOKS.....	3
1.4 JOURNALS.....	3
1.5 FIRST CONFERENCE	4
1.6 FRACTIONAL DIFFERENTIAL EQUATIONS	4
1.7 PHYSICAL SYSTEMS	5
1.8 FIRST EXPERIMENTS AND INTERPRETATIONS.....	5
1.9 DIFFERENT KINDS OF DERIVATIVES	6
1.10 USEFULNESS OF FRACTIONAL CALCULUS	6
1.11 HISTORY OF THE PROBLEM	7
1.12 OUR CONTRIBUTION	9
CHAPTER 2: PRELIMINARIES.....	12
2.1 SPACES OF INTEGRABLE FUNCTIONS AND WEIGHTED CONTINUOUS FUNCTIONS.....	13
2.2 RIEMANN-LIOUVILLE FRACTIONAL INTEGRALS AND FRACTIONAL DERIVATIVES.....	18
2.3 HADAMARD TYPE FRACTIONAL INTEGRALS AND FRACTIONAL DERIVATIVES	25
2.4 CAPUTO FRACTIONAL DERIVATIVE.....	32
2.5 SOME IMPORTANT RESULTS	32

CHAPTER 3: FRACTIONAL DIFFERENTIAL PROBLEMS WITH HILFER-RIEMANN FRACTIONAL DERIVATIVE	36
3.1 EQUIVALENCE OF THE CAUCHY TYPE PROBLEM AND THE VOLTERRA INTEGRAL EQUATION	38
3.2 EXISTENCE AND UNIQUENESS OF THE GLOBAL SOLUTION TO THE CAUCHY TYPE PROBLEM.....	44
3.3 STABILITY OF HILFER FRACTIONAL DERIVATIVE.....	52
3.4 NON-EXISTENCE OF SOLUTIONS.....	58
CHAPTER 4: FRACTIONAL DIFFERENTIAL PROBLEMS WITH HILFER-HADAMARD FRACTIONAL DERIVATIVE	68
4.1 EQUIVALENCE OF THE CAUCHY TYPE PROBLEM AND THE VOLTERRA INTEGRAL EQUATION ...	71
4.2 EXISTENCE AND UNIQUENESS OF A SOLUTION	76
4.3 STABILITY OF HILFER-HADAMARD FRACTIONAL DERIVATIVE	83
4.4 NON-EXISTENCE RESULT	92
CHAPTER 5: RECOMMENDATIONS	103
REFERENCES	105
VITA	118

THESIS ABSTRACT

Name: MOHAMMED DAHAN AHMED KASSIM

Title: WELL-POSEDNESS FOR A CAUCHY FRACTIONAL
DIFFERENTIAL PROBLEM WITH HILFER TYPE
FRACTIONAL DERIVATIVE

Degree: MASTER OF SCIENCE

Major Field: MATHEMATICS

Date of Degree: May, 2011

We considered the initial value problem for a class of nonlinear fractional differential equations that involve Hilfer and Hilfer-Hadamard fractional derivatives. The first one interpolates the Riemann-Liouville fractional derivative and the Caputo fractional derivative and the second one interpolates the Hadamard fractional derivative and its Caputo counterpart. We obtained existence and uniqueness results for both types in suitably selected underlying spaces. We proved stability results for a certain class of nonlinearity. Also, we established non-existence results in case the nonlinearity is of polynomial type using the test function method.

ملخص الرسالة

الاسم: محمد دحان أحمد قاسم

عنوان الرسالة: الصياغة الجيدة لمسألة كوشي لمعادلات تفاضلية ذات رتب غير صحيحة تحتوي على مشتقة هيلفر

الدرجة العلمية: ماجستير

التخصص: رياضيات

تاريخ التخرج: مايو 2011

درسنا مسألة القيمة الابتدائية لنوع من المعادلات التفاضلية غير الخطية ذات رتب غير صحيحة تحتوي على مشتقة هيلفر و هيلفر-هدمارد. النوع الاول يستكمل مشتقة ريمان-ليوفل ومشتقة كابوتو و النوع الثاني يستكمل مشتقة هدمارد ومشتقة كابوتو المناظرة لها. حصلنا على نتائج وجود و وحدانية الحل لكلا النوعين في فضاء تم اختياره بطريقة مناسبة. كما اثبتنا استقرار الحل لنوع معين من الدوال غير الخطية. ايضا في حالة كون الدالة الغير خطية من نوع كثيرة الحدود تم اثبات عدم وجود الحل باستخدام طريقة الدالة الاختبارية.

Chapter 1

INTRODUCTION

1.1 What is Fractional Calculus

Fractional calculus is the field of mathematical analysis which deals with the investigation and applications of integrals and derivatives of arbitrary order, unlike the classical calculus, the order of the integrals and derivatives are not necessarily of integer order. It could be any real or complex number or even a variable order.

1.2 First contributions

The idea of fractional derivative goes back to more than three hundred years. In fact, fractional calculus may be considered an old and yet novel topic. It is an old topic since, starting from some speculations of G.W. Leibniz (1697) and L. Euler (1730), it has been developed up to nowadays. A list of mathematicians, who have provided important contributions up to the middle of the last century, includes P. S. Laplace (1812), J. B. J. Fourier (1822), N. H. Abel (1823-1826), J. Liouville (1832-1873), H. Laurent (1884), P. A. Nekrassov (1888), A. Krug (1890), J. Hadamard (1892), O. Heaviside (1892-1912), S. Pincherle (1902), G. H. Hardy and J.E. Littlewood (1917-1928), H. Weyl (1917), P. Lévy (1923), A. Marchaud (1927), H. T. Davis (1924-1936), A. Zygmund (1935-1945), E. R. Love (1938-1996), A. Erdélyi (1939-1965), H. Kober (1940), D. V. Widder (1941), M. Riesz (1949),...

1.3 First books

The first book entirely devoted to the subject is the book by Oldham and Spanier [79]. Other books are by: Samko, Kilbas and Marichev [89], Podlubny [83], Kilbas, Strivastava and Trujillo [55], Mainardi [68]. For the first monograph the merit is ascribed to K. B. Oldham and J. Spanier, see [79], who, after a joint collaboration started in 1968, published a book devoted to fractional calculus in 1974. Nowadays, the list of texts and proceedings devoted solely or partly to fractional calculus and its applications includes about a dozen of titles [87, 88], among which the encyclopedic treatise by Samko, Kilbas & Marichev [89] is the most prominent.

Furthermore, we draw the attention of the reader to the treatises by Davis [17], Erdélyi [36], Gelfand and Shilov [43], Dzherbashian [25, 26], Caputo [16], Babenko [8], Gorenflo and Vessella [45], which contain a detailed analysis of some mathematical aspects and/or physical applications of fractional calculus, although without explicit mention in their titles.

1.4 Journals

There are three journals in the subject: "Journal of Fractional Calculus", "Fractional Calculus and Applied Analysis" and the recent one "Fractional Dynamic Systems". These journals publish work exclusively on fractional calculus and analysis. Special issues are also being published in other journals (like : Advances in Difference equations) from time to time.

1.5 First conferenc

Fractional calculus may also be considered as a novel topic, since only from a little more than twenty years it has been the subject of specialized conferences and treatises. For the first conference the merit is ascribed to B. Ross who organized the First Conference on Fractional Calculus and its Applications at the University of New Haven in June 1974, and edited the proceedings, see [87].

1.6 Fractional Differential Equations

Fractional differential equations have been of great interest recently because of both the intensive development of the theory of fractional calculus itself and the applications of such constructions in various sciences such as physics, mechanics, chemistry, finance, engineering, possibly including fractal phenomena, etc. For details, the reader is referred to the monographs of Lakshmikantham et al. [62], Miller and Ross [73] and Samko et al. [89], Podlubny [83], Kilbas [55] and the papers of Diethelm et al. [21, 22, 23], Gaul et al [42], Glockle and Nonnenmacher [44], Mainardi [67], Metzler et al. [71], Momani and Hadid [75], Momani et al. [76], Podlubny et al. [84], Yu and Gao [99] and the references therein.

1.7 Physical Systems

It should be noted that there are a growing number of physical systems whose behavior can be compactly described using fractional-order system theory. Of specific interest to engineers are viscoelastic materials [12, 57, 58, 92], electrochemical processes [50, 94], long lines [48], dielectric polarization [93], colored noise [69], and chaos [46]. With the growing number of applications, it is important to establish a clear initialized system theory for these fractional-order systems, so that it may be accessible to the general engineering and scientific communities.

1.8 First experiments and interpretations

Four decades ago engineers were ignoring fractional derivatives and avoided dealing with them directly because of their complexity and sometimes inconsistency. There was no solid basis as their study is very different from the integer order case. The interest to fractional calculus has been accelerated in the past three decades after the publication of the three papers of Bagley and Torvik [9-11] and the paper by Podlubny [83]. The authors proved by experiments that when using fractional derivatives for viscoelastic materials leads to:

- experimental data (amplitude/frequency and dispersions) are in agreement within a broad frequency range,
- time domain responses as stress relaxation and creep are well represented,
- the number of parameters is significantly reduced.

In addition to that, Podlubny established in [85] a geometric interpretation of the fractional integral and a physical interpretation of the fractional derivative.

1.9 Different kinds of derivatives

Several kinds of fractional derivatives have been defined: The Riemann-Liouville, Weyl, Weyl-Liouville, Chen, Bessel, Marchaud, Grunwald-Letnikov, Dzerbashyan, Hadamard derivative, etc... The first two are the most used ones. The first one is commonly used especially by mathematicians but it does not work with the usual initial conditions as the physical meaning of initial fractional derivatives is not clear and there are no ways how to measure them. Solutions must behave like the inverse of a power type function nearly the initial point. Physicists and engineers preferred the Caputo derivative which is compatible with the usual initial conditions although we can find nowadays several tentative justifying the use of fractional initial conditions.

1.10 Usefulness of fractional calculus

Fractional calculus has attracted a large number of researchers and has become very popular in numerous fields of sciences and engineering. It has been shown that fractional derivatives and fractional integrals are very successful in describing for instance anomalous kinetics and continuous time random walks. In general it has been confirmed that fractional derivatives are more suitable and adequate than derivatives of integer orders for the description of properties of many materials (especially materi-

als with memory) and hereditary phenomena and processes. There are many other complex systems in atmospheric diffusion of pollution, network traffic, anomalous diffusion, chaotic processes, biology, medicine, modeling and identification, electronics and wave propagation, mechanics, astrophysics, signal processing, chaotic dynamics, optics and porous media where fractional calculus has applications.

1.11 History of the problem

For the Riemann-Liouville Fractional Differential Problem (R-L FDP)

$$\begin{cases} (D_{a+}^{\alpha} y)(x) = f[x, y(x)], & x > a, \quad 0 < \alpha < 1 \\ (D_{a+}^{\alpha-1} y)(a+) = c, \end{cases} \quad (1.1)$$

and the Caputo Fractional Differential Problem (C FDP)

$$\begin{cases} ({}^C D_{a+}^{\alpha} y)(x) = f[x, y(x)], & x > a, \quad 0 < \alpha < 1 \\ y(a+) = c, \end{cases} \quad (1.2)$$

we may quote almost all the papers in our references.

Most of these publications are concerned with the issue of existence and uniqueness.

We can cite the first work of Pitcher and W. Sewell [82] who in 1938 studied the nonlinear fractional differential equation (1.1) when f is bounded and Lipschitz continuous with respect to y . But special cases of f have been treated before that by some researches such as O'Shaughnessy [80] (1918), Post [86] (1919) and Mandelbrojt

[70] (1925).

The R-L FDP (1.1) was studied for the first time by Al-Bassam [6] (1965) in the space of continuous functions when f is a real-valued continuous and Lipschitz in a certain domain.

Then we can find the papers of Al-Abedeen [4] and Al-Abedeen and Arora [5], Arora and Alshamani [7], Tazali [95], Tazali and Karim [96] when the nonlinearity satisfies a generalized Lipschitz condition using different methods. We also quote the works of Leskovskij [64], Semenchuk [90], Luszczki and Rzepecki [66], El-Sayed and El-Sayed and collaborators [27-35]. In almost all these publications there was a serious problem regarding the equivalence between the Fractional Differential Problem and its corresponding Integral Equation. These publications were followed by many others written by Delbosco and Rodino [18], Hayek et al. [47], Podlubny [83], Kilbas et al. [51,52], Diethelm and Ford [24], Kilbas and Marzan ([53] and [54] where the equivalence was proved rigorously, using successive expansion and fixed point theorems. In parallel we can find many works on special nonlinearities f , the linear case and also on fractional differential problems involving other types of fractional derivatives than the Riemann-Liouville fractional derivative.

More recently, in addition to the Picard-type iterative process and the approximate-iterative method by Dzjadyk, several kinds of fixed point theorems have been used in [2,14,3,53,98,65,99,97,54,59,61,13,24,78,91,100].

Some stability results have been established in [1], [34], [27-34], [56], [35], [37], [77].

Finally we mention that the quasilinearization method has been generalized to the fractional order case in [19,20].

1.12 Our contribution

In [49], Hilfer introduced a new type of fractional derivative which we name after him in this thesis

$$\left(D_{a\pm}^{\alpha,\beta}y\right)(x) = \pm \left(I_{a\pm}^{\beta(1-\alpha)} \frac{d}{dx} I_{a\pm}^{(1-\beta)(1-\alpha)} y\right)(x). \quad (1.3)$$

This fractional derivative may be viewed as interpolating the Riemann-Liouville derivative and the Caputo derivative in the sense that for $\beta = 0$ we recover the Riemann-Liouville fractional derivative (1.1) and for $\beta = 1$ we find the Caputo fractional derivative (1.2).

We intend to study the basic fractional differential problem with the fractional derivative of Hilfer type :

$$\begin{cases} \left(D_{a+}^{\alpha,\beta}y\right)(x) = f[x, y(x)], \quad x > a, \quad 0 < \alpha < 1, \quad 0 \leq \beta \leq 1 \\ \left(I_{a+}^{(1-\beta)(1-\alpha)}y\right)(a) = c. \end{cases} \quad (1.4)$$

In this thesis we are interested in problem (1.4). It had appeared for the first time in [49] but we could not find any mathematical basic treatment (well-posedness, stability, asymptotic behavior, ...) for this problem.

The topics to be covered in this thesis are the following

- Existence and uniqueness of solutions
- Some stability issues
- Non-existence and/or blow up of solutions.

As it is mentioned earlier, the equation with Riemann-Liouville derivative or Caputo

is supplemented by an initial value condition which depends on the considered type of fractional derivative. These two problems are mathematically and physically different. Nevertheless, we hope that our investigation will lead to some sort of unification of both treatments.

For the existence and uniqueness we will first establish an equivalence between the FDP (1.4) and an integral equation. This integral equation will determine the operator T for which we apply later a suitable fixed point theorem. We will follow the nice proof in the book of Kilbas et al. [55] where they first establish the equivalence between the Fractional Differential Problem and its corresponding Nonlinear Volterra Integral Equation, then using the contraction principal they establish the existence and uniqueness of a solution.

This equivalence is crucial and more important is the space where it holds true. This space is delicate and should be carefully selected as fractional derivatives (apart from Caputo fractional derivative) cause some regularity problems as they (and the functions) may be singular at the initial point.

For the stability we will exploit the “initial decay” of solutions, that is the behavior of solutions nearby the initial point and determine sufficient conditions on the nonlinearity which allow us to push and keep this behavior for all time in case of global existence. We apply some desingularization techniques to be able to use a singular version of the Gronwall inequality.

There are several ways to investigate the non-existence and the blow up of solutions. We use the test function method developed by Pohozaev and Mitidieri [74]. This technique is very efficient and it has been used successfully for the case of Riemann-

Liouville fractional derivative. It works perfectly for a polynomial source. We will follow in particular the paper by Laskri and Tatar [63].

This thesis is organized as follows:

In chapter two, we present the basic definitions, lemmas, properties and notation needed later in this thesis. In chapter three, the existence and uniqueness of a solution, a stability result and a non-existence result are stated and proved. The same kinds of questions are addressed for the corresponding Hilfer-Hadamard type problem in chapter four. Some recommendations for future work are presented in chapter five.

Chapter 2

PRELIMINARIES

In this chapter we present some definitions, lemmas, properties and notation which will be used in our theorems later.

2.1 Spaces of Integrable Functions and Weighted Continuous Functions

In this section we present the definition of p -integrable functions, weighted p -integrable functions and some weighted spaces of continuous functions. We also give some characterizations of these modified spaces which will be used later. Moreover, some important embeddings are stated.

Definition 2.1.1. [55, p. 1]: Let $\Omega = (a, b)$ ($-\infty \leq a < b \leq \infty$) be a finite or infinite interval of the real axis $\mathbb{R} = (-\infty, \infty)$. We denote by $L_p(a, b)$ ($1 \leq p \leq \infty$) the set of those Lebesgue real-valued measurable functions f on $[a, b]$ for which $\|f\|_p < \infty$, where

$$\|f\|_p = \left(\int_a^b |f(t)|^p dt \right)^{1/p}, \quad 1 \leq p < \infty,$$

and

$$\|f\|_\infty = \operatorname{ess\,sup}_{a \leq x \leq b} |f(x)|.$$

Here $\operatorname{ess\,sup} |f(x)|$ is the essential supremum of the function $|f(x)|$.

Definition 2.1.2. [55, p. 1]: The space $X_c^p(a, b)$ ($c \in \mathbb{R}$, $1 \leq p \leq \infty$) consists of those real-valued Lebesgue measurable functions g on (a, b) for which $\|g\|_{X_c^p} < \infty$,

where

$$\|g\|_{X_c^p} = \left(\int_a^b |t^c g(t)|^p \frac{dt}{t} \right)^{1/p}, \quad 1 \leq p < \infty, \quad c \in \mathbb{R}$$

and

$$\|g\|_{X_c^\infty} = \operatorname{ess\,sup}_{a \leq x \leq b} |x^c g(x)|, \quad c \in \mathbb{R}.$$

In particular, when $c = 1/p$ we see that $X_{1/p}^p(a, b) = L_p(a, b)$.

Definition 2.1.3. [55, p. 3]: Let $\Omega = [a, b]$ ($-\infty < a < b < \infty$) and $m \in \mathbb{N}_0 = \{0, 1, \dots\}$. We denote by $C^m(\Omega)$ the space of functions f which are m times continuously differentiable on Ω with the norm

$$\|f\|_{\mathbf{C}^m} = \sum_{k=0}^m \|f^{(k)}\|_{\mathbf{C}} = \sum_{k=0}^m \max_{x \in \Omega} |f^{(k)}(x)|, \quad m \in \mathbb{N}_0.$$

In particular, for $m = 0$, $C^0(\Omega) \equiv C(\Omega)$ is the space of continuous function f on Ω with the norm

$$\|f\|_{\mathbf{C}} = \max_{x \in \Omega} |f(x)|.$$

Definition 2.1.4. [55, p. 4]: Let $\Omega = [a, b]$ be a finite interval and $0 \leq \gamma < 1$, we introduce the weighted space $C_\gamma[a, b]$ of continuous functions f on $(a, b]$

$$C_\gamma[a, b] = \{f : (a, b] \rightarrow \mathbb{R} : (x - a)^\gamma f(x) \in C[a, b]\}.$$

In the space $C_\gamma[a, b]$, we define the norm

$$\|f\|_{C_\gamma} = \|(x - a)^\gamma f(x)\|_C, \quad C_0[a, b] = C[a, b].$$

Definition 2.1.5. [55, p. 4]: For $n \in \mathbb{N}$ we denote by $C_\gamma^n[a, b]$ ($0 \leq \gamma < 1$) the Banach space of functions f which are continuously differentiable on $[a, b]$ up to order $n - 1$ and have the derivative $f^{(n)}$ of order n on $(a, b]$ such that $f^{(n)} \in C_\gamma[a, b]$

$$C_\gamma^n[a, b] = \{f \in C^{n-1}[a, b] : f^{(n)} \in C_\gamma[a, b]\},$$

with the norm

$$\|f\|_{C_\gamma^n} = \sum_{k=0}^{n-1} \|f^{(k)}\|_C + \|f^{(n)}\|_{C_\gamma}, \quad C_\gamma^0[a, b] = C_\gamma[a, b].$$

From this definition we have the following characterization of the space $C_\gamma^n[a, b]$.

Lemma 2.1.1. [55, p. 4]: Let $n \in \mathbb{N} = \{1, 2, \dots\}$ and $0 \leq \gamma < 1$. The space $C_\gamma^n[a, b]$ consists of those and only those functions f which can be represented in the form

$$f(x) = \frac{1}{(n-1)!} \int_a^x (x-t)^{n-1} \varphi(t) dt + \sum_{k=0}^{n-1} c_k (x-a)^k,$$

where $\varphi \in C_\gamma[a, b]$ and c_k ($k = 0, 1, \dots, n-1$) are arbitrary constants.

Moreover,

$$\varphi(t) = f^{(n)}(t), \quad c_k = \frac{f^{(k)}(a)}{k!} \quad (k = 0, 1, \dots, n-1).$$

Definition 2.1.6. [55, p. 4]: Let $\Omega = [a, b]$ ($0 < a < b < \infty$) be a finite interval and $0 \leq \gamma < 1$, we introduce the weighted space $C_{\gamma, \log}[a, b]$ of continuous functions g

on $(a, b]$

$$C_{\gamma, \log} [a, b] = \left\{ g : (a, b] \rightarrow \mathbb{R} : \left(\log \frac{x}{a} \right)^\gamma g(x) \in C[a, b] \right\}.$$

In the space $C_{\gamma, \log} [a, b]$, we define the norm

$$\|g\|_{C_{\gamma, \log}} = \left\| \left(\log \frac{x}{a} \right)^\gamma g(x) \right\|_C, \quad C_{0, \log} [a, b] = C[a, b].$$

Definition 2.1.7. [55, p. 5]: Let $\delta = x \frac{d}{dx}$ be the δ -derivative, for $n \in \mathbb{N}$ we denote by $C_{\delta, \gamma}^n [a, b]$ ($0 \leq \gamma < 1$) the Banach space of functions g which have continuous δ -derivatives on $[a, b]$ up to order $n - 1$ and have the derivative $\delta^n g$ of order n on $(a, b]$ such that $\delta^n g \in C_{\gamma, \log} [a, b]$

$$C_{\delta, \gamma}^n [a, b] = \left\{ \delta^k g \in C[a, b], \quad k = 0, \dots, n - 1, \quad \delta^n g \in C_{\gamma, \log} [a, b] \right\}$$

with the norm

$$\|g\|_{C_{\delta, \gamma}^n} = \sum_{k=0}^{n-1} \|\delta^k g\|_C + \|\delta^n g\|_{C_{\gamma, \log}}.$$

When $n = 0$ we set

$$C_{\delta, \gamma}^0 [a, b] = C_{\gamma, \log} [a, b].$$

From this definition we have the following characterization of the space $C_{\delta, \gamma}^n [a, b]$.

Lemma 2.1.2. [55, p. 5]: Let $0 < a < b < \infty$, $n \in \mathbb{N}_0 = \{0, 1, \dots\}$ and $0 \leq \gamma < 1$. The space $C_{\delta, \gamma}^n [a, b]$ consists of those and only those functions g which can be

represented in the form

$$g(x) = \frac{1}{(n-1)!} \int_a^x \left(\log \frac{x}{t} \right)^{n-1} \varphi(t) \frac{dt}{t} + \sum_{k=0}^{n-1} d_k \left(\log \frac{x}{a} \right)^k,$$

where $\varphi \in C_{\gamma, \log}[a, b]$ and d_k ($k = 0, 1, \dots, n-1$) are arbitrary constants.

Moreover,

$$\varphi(t) = (\delta^n g)(t), \quad d_k = \frac{(\delta^k g)(a)}{k!} \quad (k = 0, 1, \dots, n-1).$$

From the Definitions 2.1.3 to 2.1.7 we have the following embeddings.

Property 2.1.1. [55, p. 5]: Let $n \in \mathbb{N}_0 = \{0, 1, \dots\}$ and let μ_1 and μ_2 be real numbers such that

$$0 \leq \mu_1 \leq \mu_2 < 1.$$

The following embedding hold:

$$C^n[a, b] \rightarrow C_{\mu_1}^n[a, b] \rightarrow C_{\mu_2}^n[a, b],$$

with

$$\|f\|_{C_{\mu_2}^n} \leq K \|f\|_{C_{\mu_1}^n}, \quad K = \min \left[1, (b-a)^{\mu_2-\mu_1} \right];$$

$$C_{\delta}^n[a, b] \rightarrow C_{\delta, \mu_1}^n[a, b] \rightarrow C_{\delta, \mu_2}^n[a, b],$$

with

$$\|f\|_{C_{\delta, \mu_2}^n} \leq K_{\delta} \|f\|_{C_{\delta, \mu_1}^n}, \quad K_{\delta} = \min \left[1, \left(\log \frac{b}{a} \right)^{\mu_2-\mu_1} \right].$$

In particular,

$$C[a, b] \rightarrow C_{\mu_1}[a, b] \rightarrow C_{\mu_2}[a, b],$$

with

$$\|f\|_{C_{\mu_2}} \leq (b-a)^{\mu_2-\mu_1} \|f\|_{C_{\mu_1}};$$

$$C[a, b] \rightarrow C_{\mu_1, \log}[a, b] \rightarrow C_{\mu_2, \log}[a, b],$$

with

$$\|f\|_{C_{\mu_2, \log}} \leq \left(\log \frac{b}{a}\right)^{\mu_2-\mu_1} \|f\|_{C_{\mu_1, \log}}.$$

Definition 2.1.8. [55, p. 24]: *The Euler gamma function $\Gamma(z)$ is defined by the so-called Euler integral of the second kind:*

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad z > 0,$$

where $t^{z-1} = e^{(z-1) \log t}$.

2.2 Riemann-Liouville Fractional Integrals and Fractional Derivatives

In this section we introduce the definitions of the Riemann-Liouville fractional integrals and fractional derivatives on a finite interval of the real line and present some of their properties in the space of continuous functions.

Definition 2.2.1. [55, p. 69]: *Let $\Omega = [a, b]$ ($-\infty < a < b < \infty$) be a finite*

interval on the real axis \mathbb{R} . The Riemann-Liouville left-sided fractional integral $I_{a+}^\alpha f$ of order $\alpha > 0$ is defined by

$$(I_{a+}^\alpha f)(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \frac{f(t)}{(x-t)^{1-\alpha}} dt, \quad (a < x < b, \alpha > 0)$$

provided that the integral exists. Here $\Gamma(\alpha)$ is the Gamma function. When $\alpha = 0$, we set

$$I_{a+}^0 f = f.$$

Definition 2.2.2. [55, p. 69]: Let $\Omega = [a, b]$ ($-\infty < a < b < \infty$) be a finite interval on the real axis \mathbb{R} . The Riemann-Liouville right-sided fractional integral $I_{b-}^\alpha f$ of order $\alpha > 0$ is defined by

$$(I_{b-}^\alpha f)(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \frac{f(t)}{(t-x)^{1-\alpha}} dt, \quad (a \leq x < b, \alpha > 0)$$

provided that the integral exists. When $\alpha = 0$, we define

$$I_{b-}^0 f = f.$$

Definition 2.2.3. [55, p. 70]: Let $\Omega = [a, b]$ ($-\infty < a < b < \infty$) be a finite interval on the real axis \mathbb{R} . The Riemann-Liouville left-sided fractional derivative $D_{a+}^\alpha f$ of order α ($0 \leq \alpha < 1$) is defined by

$$(D_{a+}^\alpha f)(x) = \frac{d}{dx} (I_{a+}^{1-\alpha} f)(x)$$

that is

$$(D_{a+}^{\alpha} f) = \frac{1}{\Gamma(1-\alpha)} \frac{d}{dx} \int_a^x \frac{f(t)}{(x-t)^{\alpha}} dt \quad (x > a, \quad 0 < \alpha < 1),$$

when $\alpha = 1$ we have $D_{a+}^{\alpha} f = Df$. In particular, when $\alpha = 0$, then

$$D_{a+}^0 f = f.$$

Definition 2.2.4. [55, p. 70]: Let $\Omega = [a, b]$ ($-\infty < a < b < \infty$) be a finite interval on the real axis \mathbb{R} . The Riemann-Liouville right-sided fractional derivative $D_{b-}^{\alpha} f$ of order α ($0 \leq \alpha < 1$) is defined by

$$(D_{b-}^{\alpha} f)(x) = -\frac{d}{dx} (I_{b-}^{1-\alpha} f)(x)$$

that is

$$(D_{b-}^{\alpha} f) = -\frac{1}{\Gamma(1-\alpha)} \frac{d}{dx} \int_x^b \frac{f(t)}{(t-x)^{\alpha}} dt \quad (a \leq x < b, \quad 0 < \alpha < 1).$$

In particular, when $\alpha = 0$, then

$$D_{b-}^0 f = f.$$

It can be directly verified that the Riemann-Liouville fractional integral and fractional derivative of the power function $(x-a)^{\beta-1}$ ($\beta > 0$) yield the same power function with α added or subtracted from the power $\beta > 0$ with a certain coefficient in front of this power function.

Property 2.2.1. [55, p. 71]: If $\alpha \geq 0$ and $\beta > 0$, then

$$\left(I_{a+}^{\alpha} (t-a)^{\beta-1}\right)(x) = \frac{\Gamma(\beta)}{\Gamma(\beta+\alpha)} (x-a)^{\beta+\alpha-1}, \quad \alpha \geq 0$$

$$\left(D_{a+}^{\alpha} (t-a)^{\beta-1}\right)(x) = \frac{\Gamma(\beta)}{\Gamma(\beta-\alpha)} (x-a)^{\beta-\alpha-1}, \quad \alpha \geq 0.$$

In particular, if $\beta = 1$ and $\alpha \geq 0$, then the Riemann-Liouville fractional derivative of a constant is not equal to zero:

$$\left(D_{a+}^{\alpha} 1\right)(x) = \frac{(x-a)^{-\alpha}}{\Gamma(1-\alpha)}, \quad \alpha \geq 0.$$

Property 2.2.2. [55, p. 72]: If $0 < \alpha \leq 1$, then

$$\left(D_{a+}^{\alpha} (t-a)^{\alpha-1}\right)(x) = 0.$$

Lemma 2.2.1. [60, p. 6]: If $0 \leq \gamma < 1$, then the fractional integration operator

I_{a+}^{α} of order α ($\alpha > 0$) is bounded in $C_{\gamma}[a, b]$:

$$\|I_{a+}^{\alpha} g\|_{C_{\gamma}} \leq \frac{\Gamma(1-\gamma)}{\Gamma(1+\alpha-\gamma)} (b-a)^{\alpha} \|g\|_{C_{\gamma}},$$

here I_{a+}^{α} is the Riemann-Liouville fractional integral operator and $g \in C_{\gamma}[a, b]$.

Furthermore, we have the following important lemmas.

Lemma 2.2.2. [60, p. 6]: *The fractional integration operator I_{a+}^α of order α ($\alpha > 0$) is a mapping from $C[a, b]$ to $C[a, b]$, and*

$$\|I_{a+}^\alpha g\|_C \leq \frac{(b-a)^\alpha}{\alpha \Gamma(\alpha)} \|g\|_C,$$

where $g \in C[a, b]$.

Lemma 2.2.3. [60, p. 5]: *Let $0 \leq \gamma < 1$, $a < c < b$, $g \in C_\gamma[a, c]$ and $g \in C[c, b]$. Then $g \in C_\gamma[a, b]$ and*

$$\|g\|_{C_\gamma[a, b]} \leq \max \left[\|g\|_{C_\gamma[a, c]}, (b-a)^\gamma \|g\|_{C[c, b]} \right].$$

Now we consider some other properties of the Riemann-Liouville fractional integral (Definition 2.2.1) and the Riemann-Liouville fractional derivative (Definition 2.2.3) in the space $C_\gamma[a, b]$ and $C_\gamma^n[a, b]$ defined in Definition 2.1.4 and Definition 2.1.5, respectively. The existence of the fractional integral $I_{a+}^\alpha f$ in the space $C_\gamma[a, b]$ and the fractional derivative $D_{a+}^\alpha f$ in the space $C_\gamma^n[a, b]$ are given by the following lemmas.

Lemma 2.2.4. [55, p. 76]: *The following hold*

(a) *Let $\alpha > 0$ and $0 \leq \gamma < 1$.*

If $\gamma > \alpha$, then the fractional integration operator I_{a+}^α is bounded from $C_\gamma[a, b]$ into $C_{\gamma-\alpha}[a, b]$:

$$\|I_{a+}^\alpha f\|_{C_{\gamma-\alpha}} \leq k_1 \|f\|_{C_\gamma},$$

$$k_1 = \frac{\Gamma(1-\gamma)}{\Gamma(1+\alpha-\gamma)}.$$

In particular I_{a+}^α is bounded in $C_\gamma[a, b]$.

(b) If $\gamma \leq \alpha$, then the fractional integration operator I_{a+}^α is bounded from $C_\gamma[a, b]$ into $C[a, b]$:

$$\|I_{a+}^\alpha f\|_C \leq k_2 \|f\|_{C_\gamma},$$

$$k_2 = (b-a)^{\alpha-\gamma} \frac{\Gamma(1-\gamma)}{\Gamma(1+\alpha-\gamma)}.$$

In particular I_{a+}^α is bounded in $C_\gamma[a, b]$.

Lemma 2.2.5. [55, p. 77]: Let $0 < \alpha < 1$ and $0 \leq \gamma < 1$. If $f \in C_\gamma^1$, then the fractional derivatives D_{a+}^α and D_{b-}^α exist on $(a, b]$ and $[a, b)$ respectively, and can be represented in the forms

$$(D_{a+}^\alpha f)(x) = \frac{1}{\Gamma(1-\alpha)} \left[\frac{f(a)}{(x-a)^\alpha} + \int_a^x \frac{f'(t) dt}{(x-t)^\alpha} \right]$$

and

$$(D_{b-}^\alpha f)(x) = \frac{1}{\Gamma(1-\alpha)} \left[\frac{f(b)}{(b-x)^\alpha} - \int_x^b \frac{f'(t) dt}{(t-x)^\alpha} \right],$$

Lemma 2.2.6. [55, p. 77]: (The Semigroup Property of the fractional integration operator I_{a+}^α) Let $\alpha > 0$, $\beta > 0$ and $0 \leq \gamma < 1$. If $f \in C_\gamma[a, b]$ then the equation

$$I_{a+}^\alpha I_{a+}^\beta f = I_{a+}^{\alpha+\beta} f$$

holds at any point $x \in (a, b]$. When $f \in C[a, b]$ this relation is valid at any point $x \in [a, b]$.

The following assertion shows that the fractional differentiation is an inverse operation

to the fractional integration from the left.

Lemma 2.2.7. [55, p. 77]: *Let $\alpha > 0$ and $0 \leq \gamma < 1$. If $f \in C_\gamma[a, b]$, then the relation*

$$D_{a+}^\alpha I_{a+}^\alpha f = f$$

holds at any point $x \in (a, b]$. When $f \in C[a, b]$ this relation is valid at any point $x \in [a, b]$.

Another composition property between the fractional differentiation operator (Definition 2.2.3) and the fractional integration (Definition 2.2.1) is given next.

Property 2.2.3. [55, p. 77]: *Let $\alpha > \beta > 0$ and $0 \leq \gamma < 1$. If $f \in C_\gamma[a, b]$, then the relation*

$$D_{a+}^\beta I_{a+}^\alpha f = I_{a+}^{\alpha-\beta} f$$

holds at any point $x \in (a, b]$. When $f \in C[a, b]$ this relation is valid at any point $x \in [a, b]$. In particular, when $\beta = k \in \mathbb{N}$ and $\alpha > k$, then $D_{a+}^k I_{a+}^\alpha f = I_{a+}^{\alpha-k} f$.

The following result provides another composition of the fractional integration operator I_{a+}^α with the fractional differentiation operator D_{a+}^α .

Lemma 2.2.8. [55, p. 77]: *Let $0 < \alpha < 1$, $0 \leq \gamma < 1$. Also let $I_{a+}^{1-\alpha} f$ be the fractional integral of the function f of order $1 - \alpha$.*

If $f \in C_\gamma[a, b]$ and $I_{a+}^{1-\alpha} f \in C_\gamma^1[a, b]$, then the equality

$$(I_{a+}^\alpha D_{a+}^\alpha f)(x) = f(x) - \frac{(I_{a+}^{1-\alpha} f)(a)}{\Gamma(\alpha)} (x - a)^{\alpha-1},$$

holds at any point $x \in (a, b]$.

Lemma 2.2.9. [55, p. 76]: (Fractional Integration by Parts) : *Let $\alpha > 0$, $p \geq 1$, $q \geq 1$ and $\frac{1}{p} + \frac{1}{q} \leq 1 + \alpha$ ($p \neq 1$ and $q \neq 1$ in the case when $\frac{1}{p} + \frac{1}{q} = 1 + \alpha$). If $\varphi \in L_p(a, b)$ and $\psi \in L_q(a, b)$, then*

$$\int_a^b \varphi(x) (I_{a+}^\alpha \psi)(x) dx = \int_a^b \psi(x) (I_{b-}^\alpha \varphi)(x) dx.$$

2.3 Hadamard Type Fractional Integrals and Fractional Derivatives

In this section we present the definitions and some properties of the Hadamard type fractional integrals and fractional derivatives which will be involved in the problems investigated later.

Definition 2.3.1. [55, p. 110]: *Let (a, b) ($0 \leq a < b \leq \infty$) be a finite or infinite interval of the half-axis \mathbb{R}^+ and let $\alpha > 0$. The Hadamard left-sided fractional integral $\mathcal{J}_{a+}^\alpha f$ of order $\alpha > 0$ is defined by*

$$(\mathcal{J}_{a+}^\alpha f)(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f(t) dt}{t} \quad (a < x < b)$$

provided that the integral exists. When $\alpha = 0$, we set

$$\mathcal{J}_{a+}^0 f = f.$$

Definition 2.3.2. [55, p. 110]: Let (a, b) ($0 \leq a < b \leq \infty$) be a finite or infinite interval of the half-axis \mathbb{R}^+ and let $\alpha > 0$. The Hadamard right-sided fractional integral $\mathcal{J}_{b-}^\alpha f$ of order $\alpha > 0$ is defined by

$$(\mathcal{J}_{b-}^\alpha f)(x) := \frac{1}{\Gamma(\alpha)} \int_x^b \left(\log \frac{t}{x} \right)^{\alpha-1} \frac{f(t) dt}{t} \quad (a < x < b)$$

provided that the integral exists. When $\alpha = 0$, we set

$$\mathcal{J}_{b-}^0 f = f.$$

Definition 2.3.3. [55, p. 111]: The left-sided Hadamard fractional derivative of order $0 \leq \alpha < 1$ on (a, b) is defined by

$$(\mathcal{D}_{a+}^\alpha f)(x) := \delta \left(\mathcal{J}_{a+}^{1-\alpha} f \right) (x),$$

that is

$$(\mathcal{D}_{a+}^\alpha f)(x) = \left(x \frac{d}{dx} \right) \frac{1}{\Gamma(1-\alpha)} \int_a^x \left(\log \frac{x}{t} \right)^{-\alpha} \frac{f(t) dt}{t} \quad (a < x < b).$$

In particular, when $\alpha = 0$ we have

$$\mathcal{D}_{a+}^0 f = f.$$

Definition 2.3.4. [55, p. 111]: *The right-sided Hadamard fractional derivative of order α ($0 \leq \alpha < 1$) on (a, b) is defined by*

$$(\mathcal{D}_{b-}^{\alpha} f)(x) := -\delta(\mathcal{J}_{b-}^{1-\alpha} f)(x),$$

that is

$$(\mathcal{D}_{b-}^{\alpha} f)(x) = -\left(x \frac{d}{dx}\right) \frac{1}{\Gamma(1-\alpha)} \int_x^b \left(\log \frac{t}{x}\right)^{-\alpha} \frac{f(t) dt}{t} \quad (a < x < b).$$

In particular, when $\alpha = 0$ we have

$$\mathcal{D}_{b-}^0 f = f.$$

It can be directly verified that the Hadamard fractional integral and fractional derivative of the logarithmic function $\left(\log \frac{x}{a}\right)^{\beta-1}$ yield the same logarithmic function with α added or subtracted from the power with a certain coefficient in front of the logarithmic function.

Property 2.3.1. [55, p. 112]: *If $\alpha > 0$, $\beta > 0$ and $0 < a < b < \infty$, then*

$$\left(\mathcal{J}_{a+}^{\alpha} \left(\log \frac{t}{a}\right)^{\beta-1}\right)(x) = \frac{\Gamma(\beta)}{\Gamma(\alpha + \beta)} \left(\log \frac{x}{a}\right)^{\beta+\alpha-1},$$

and

$$\left(\mathcal{D}_{a+}^{\alpha} \left(\log \frac{t}{a}\right)^{\beta-1}\right)(x) = \frac{\Gamma(\beta)}{\Gamma(\beta - \alpha)} \left(\log \frac{x}{a}\right)^{\beta-\alpha-1}.$$

In particular, if $\beta = 1$ and $\alpha \geq 0$, then the Hadamard fractional derivative of a constant is not equal to zero:

$$(\mathcal{D}_{a+}^{\alpha} 1)(x) = \frac{1}{\Gamma(1-\alpha)} \left(\log \frac{x}{a} \right)^{-\alpha},$$

when $0 < \alpha < 1$.

Lemma 2.3.1. [55, p. 113]: If $0 < \alpha \leq 1$ and $0 < a < b < \infty$. The equality $(\mathcal{D}_{a+}^{\alpha} y)(x) = 0$ is valid

$$y(x) = c \left(\log \frac{x}{a} \right)^{\alpha-1}$$

for any $c \in \mathbb{R}$.

The Hadamard fractional integra (Definition 2.3.1) satisfies the following semigroup property.

Property 2.3.2. [55, p. 118]: Let $\alpha > 0$, $\beta > 0$ and $0 \leq \mu < 1$. If $0 < a < b < \infty$, then, for $f \in C_{\mu, \log}[a, b]$

$$\mathcal{J}_{a+}^{\alpha} \mathcal{J}_{a+}^{\beta} f = \mathcal{J}_{a+}^{\alpha+\beta} f$$

holds at any point $x \in (a, b]$. When $f \in C[a, b]$ this relation is valid at any point $x \in [a, b]$.

Compositions between the fractional differentiation and fractional integration operators are given in the following properties.

Property 2.3.3. [55, p. 118]: Let $\alpha > \beta > 0$. If $f \in C_{\mu, \log}[a, b]$ and $0 < a < b < \infty$ then

$$\mathcal{D}_{a+}^{\beta} \mathcal{J}_{a+}^{\alpha} f = \mathcal{J}_{a+}^{\alpha-\beta} f$$

holds at any point $x \in (a, b]$. When $f \in C[a, b]$ this relation is valid at any point $x \in [a, b]$.

In particular, when $\beta = k \in \mathbb{N}$ and $\alpha > k$, then

$$\mathcal{D}_{a+}^k \mathcal{J}_{a+}^\alpha f = \mathcal{J}_{a+}^{\alpha-k} f.$$

The Hadamard fractional derivative is the left inverse operator to the corresponding fractional integral.

Property 2.3.4. [55, p. 116]: Let $\alpha > 0$. If $0 < a < b < \infty$ and $f \in C_{\mu, \log}[a, b]$ then

$$\mathcal{D}_{a+}^\alpha \mathcal{J}_{a+}^\alpha f = f, \quad x \in (a, b].$$

The following result provides a formula for the composition of the fractional differentiation operator \mathcal{D}_{a+}^α with the fractional integration operator \mathcal{J}_{a+}^α . It shows that fractional differentiation is not the right inverse operator of the fractional integral in general.

Theorem 2.3.1. [55, p. 119]: Let $0 < \alpha < 1$ and $0 < a < b < \infty$. Also let $\mathcal{J}_{a+}^{1-\alpha} f$ be the Hadamard fractional integral of the order $1 - \alpha$ of the function f .

If $f \in C_{\mu, \log}[a, b]$ and $\mathcal{J}_{a+}^{1-\alpha} f \in C_{\delta, \mu}^1[a, b]$, then

$$(\mathcal{J}_{a+}^\alpha \mathcal{D}_{a+}^\alpha f)(x) = f(x) - \frac{(\mathcal{J}_{a+}^{1-\alpha} f)(a)}{\Gamma(\alpha)} \left(\log \frac{x}{a} \right)^{\alpha-1}.$$

holds at any point $x \in (a, b]$. If $f \in C[a, b]$ and $\mathcal{J}_{a+}^{1-\alpha} f \in C_\delta^1[a, b]$, then the relation holds at any point $x \in [a, b]$.

Now we consider the properties of the Hadamard fractional integral and the Hadamard fractional derivative in the spaces $C_{\mu,\log}[a, b]$ and $C_{\delta,\mu}^1[a, b]$ (see Definitions 2.1.6 and 2.1.7). The existence of the fractional integral $\mathcal{J}_{a+}^\alpha f$ in the space $C_{\mu,\log}[a, b]$ and of the fractional derivative $\mathcal{D}_{a+}^\alpha y$ in the space $C_{\delta,\mu}^1[a, b]$ are given by the following assertions.

Lemma 2.3.2. [55, p. 118]: *Let $0 < a < b < \infty$, $\alpha > 0$ and $0 \leq \mu < 1$.*

(a) *If $\mu > \alpha > 0$, then the fractional integration operator \mathcal{J}_{a+}^α is bounded from $C_{\mu,\log}[a, b]$ into $C_{\mu-\alpha,\log}[a, b]$:*

$$\|\mathcal{J}_{a+}^\alpha f\|_{C_{\mu-\alpha,\log}} \leq k_1 \|f\|_{C_{\mu,\log}}$$

where

$$k_1 = \left(\log \frac{b}{a} \right)^\alpha \frac{\Gamma(1-\mu)}{\Gamma[1+\alpha-\mu]}.$$

In particular, \mathcal{J}_{a+}^α is bounded in $C_{\mu,\log}[a, b]$.

(b) *If $\mu \leq \alpha$, then the fractional integration operator \mathcal{J}_{a+}^α is bounded from $C_{\mu,\log}[a, b]$ into $C[a, b]$:*

$$\|\mathcal{J}_{a+}^\alpha f\|_C \leq k_2 \|f\|_{C_{\mu,\log}}$$

where

$$k_2 = \left(\log \frac{b}{a} \right)^{\alpha-\mu} \frac{\Gamma(1-\mu)}{\Gamma(1+\alpha-\mu)}.$$

In particular, \mathcal{J}_{a+}^α is bounded in $C_{\mu,\log}[a, b]$.

Lemma 2.3.3. [55, p. 118]: *Let $0 \leq \alpha < 1$ and $0 \leq \gamma < 1$. If $f \in C_{\gamma,\log}^1[a, b]$, then the fractional derivatives \mathcal{D}_{a+}^α and \mathcal{D}_{b-}^α exist on $(a, b]$ and $[a, b)$, respectively ($a > 0$)*

and can be represented in the forms

$$(\mathcal{D}_{a^+}^\alpha f)(x) = \frac{f(a)}{\Gamma(1-\alpha)} \left(\log \frac{x}{a}\right)^{-\alpha} + \frac{1}{\Gamma(1-\alpha)} \int_a^x \left(\log \frac{x}{t}\right)^{-\alpha} f'(t) dt,$$

and

$$(\mathcal{D}_{b^-}^\alpha f)(x) = \frac{f(b)}{\Gamma(1-\alpha)} \left(\log \frac{b}{x}\right)^{-\alpha} - \frac{1}{\Gamma(1-\alpha)} \int_x^b \left(\log \frac{t}{x}\right)^{-\alpha} f'(t) dt,$$

respectively.

Lemma 2.3.4. [55, p. 118]: *The Hadamard fractional integration operator $\mathcal{J}_{a^+}^\alpha$ of order α ($\alpha > 0$) is a mapping from $C[a, b]$ to $C[a, b]$ and*

$$\|\mathcal{J}_{a^+}^\alpha g\|_{C[a,b]} \leq \frac{(\log \frac{b}{a})^\alpha}{\alpha \Gamma(\alpha)} \|g\|_{C[a,b]},$$

where $g \in C[a, b]$.

Lemma 2.3.5. [60, p. 7]: *Let $0 \leq \gamma < 1$, $0 < a < c < b < \infty$, $g \in C_{\gamma, \log}[a, c]$ and $g \in C[c, b]$. Then $g \in C_{\gamma, \log}[a, b]$ and*

$$\|g\|_{C_{\gamma, \log}[a,b]} \leq \max \left\{ \|g\|_{C_{\gamma, \log}[a,c]}, \left(\log \frac{b}{a}\right)^\gamma \|g\|_{C[c,b]} \right\}.$$

Lemma 2.3.6. [15, p. 13] (Fractional Integration by Parts): *Let $\alpha > 0$ and $1 \leq p \leq \infty$. If $\varphi \in L_p(\mathbb{R}^+)$ and $\psi \in X_{-1/p}^q$, then*

$$\int_0^\infty \varphi(x) (\mathcal{J}_+^\alpha \psi)(x) \frac{dx}{x} = \int_0^\infty \psi(x) (\mathcal{J}_-^\alpha \varphi)(x) \frac{dx}{x},$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

2.4 Caputo Fractional Derivative

In this section we present the definitions of the Caputo fractional derivative ${}^cD_{a+}^\alpha f$ and Hadamard–Caputo fractional derivative ${}^c\mathcal{D}_{a+}^\alpha f$ of order α ($0 < \alpha < 1$), respectively.

Definition 2.4.1: Let $[a, b]$ be a finite interval of the real line \mathbb{R} . The fractional derivative ${}^cD_{a+}^\alpha f$ of order α ($0 < \alpha < 1$) on $[a, b]$ defined by

$${}^cD_{a+}^\alpha f = I_{a+}^{1-\alpha} Df$$

where $D = \frac{d}{dx}$, is called the Caputo fractional derivative of f of order α ($0 < \alpha < 1$).

Definition 2.4.2: Let (a, b) be a finite interval of the half-axis \mathbb{R}^+ . The fractional derivative ${}^c\mathcal{D}_{a+}^\alpha f$ of order α ($0 < \alpha < 1$) on (a, b) defined by

$${}^c\mathcal{D}_{a+}^\alpha f = \mathcal{J}_{a+}^{1-\alpha} \delta f$$

where $\delta = x \frac{d}{dx}$, is called the Hadamard–Caputo fractional derivative of order α .

2.5 Some Important Results

In this section we present some other important definitions, lemmas, theorems and properties. These will determine the assumption, tools and the methods utilized in our results later.

Definition 2.5.1. [60, p. 3]: (*Lipschitz condition*) Assume that $f[., y(.)]$ is defined on the set $(a, b] \times \mathbf{G}$ ($\mathbf{G} \subset \mathbb{R}$). $f[., y(.)]$ is said to satisfy the Lipschitz condition with respect to the second variable, if for all $x \in (a, b]$ and for any $y_1, y_2 \in \mathbf{G} \subset \mathbb{R}$ one has

$$|f[x, y_1] - f[x, y_2]| \leq A |y_1 - y_2|,$$

where $A > 0$ does not depend on $x \in (a, b]$. In this case we say that the function is Lipschitz continuous with respect to y with Lipschitz constant A .

Lemma 2.5.1. [72, p. 23]: If $\lambda, \nu, \omega > 0$, then for any $t > 0$, we have

$$\int_0^t (t-s)^{\nu-1} s^{\lambda-1} e^{-\omega s} ds \leq C t^{\nu-1},$$

where C is a positive constant independent of t . In fact,

$$C = \max \{1, 2^{1-\nu}\} \Gamma(\lambda) (1 + \lambda(\lambda+1)/\nu) \omega^{-\lambda}.$$

Lemma 2.5.2. [41, p. 581]: We have, for positive a, b and $\gamma > 1$, the inequality

$$(a+b)^\gamma \leq 2^{\gamma-1} (a^\gamma + b^\gamma).$$

Lemma 2.5.3. [81, p. 126]: Let a, b be two continuous, positive functions defined on $[t_0, \infty)$, $t_0 \geq 0$, and $w : [0, \infty) \rightarrow [0, \infty)$ be a continuous monotonic nondecreasing function such that $w(0) = 0$ and $w(x) > 0$ for $x > 0$. If u is a positive differentiable

function on $[t_0, \infty)$ that satisfies

$$u'(t) \leq a(t) w(u(t)) + b(t), \quad t \in [t_0, \infty),$$

then we have

$$u(t) \leq G^{-1} \left[G \left(u(t_0) + \int_{t_0}^t b(s) ds \right) + \int_{t_0}^t a(s) ds \right],$$

for the values of t , for which the right-hand side is well-defined, where

$$G(r) = \int_{r_0}^r \frac{ds}{w(s)}, \quad r > r_0 > 0.$$

Theorem 2.5.1. [55, p. 68]: (*Banach Fixed Point Theorem*) Let (U, d) be a nonempty complete metric space, let $0 \leq w < 1$, and let $T : U \rightarrow U$ be a map such that, for every $u, v \in U$, the relation

$$d(Tu, Tv) \leq wd(u, v), \quad 0 \leq w < 1$$

holds. Then the operator T has a unique fixed point $u^* \in U$.

Furthermore, if T^k ($k \in \mathbb{N}$) is the sequence of operators defined by

$$T^1 = T \text{ and } T^k = TT^{k-1} \text{ } (k \in \mathbb{N} \setminus \{1\}),$$

then, for any $u_0 \in U$, the sequence $\{T^k u_0\}_{k=1}^\infty$ converges to the above fixed point u^* .

Theorem 2.5.2. [63, p. 1268] : (Young's inequality) *If a and b are nonnegative real numbers and p and q are positive real numbers such that $1/p + 1/q = 1$ then we have*

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

Equality holds if and only if $a^p = b^q$.

Chapter 3

Fractional Differential Problems

with Hilfer-Riemann Fractional

Derivative

This chapter is devoted to proving the existence and uniqueness of solutions to a Cauchy type problem of fractional order on a finite interval of the real axis in a weighted space of continuous functions.

Also a stability result and a non-existence result are stated and proved with Hilfer type fractional derivative.

We consider the Cauchy type problem

$$\begin{cases} \left(D_{a+}^{\alpha,\beta} y \right) (x) = f [x, y (x)], & x > a, \quad 0 < \alpha < 1, \quad 0 \leq \beta \leq 1 \\ \left(I_{a+}^{(1-\beta)(1-\alpha)} y \right) (a) = c. \end{cases} \quad (3.1)$$

in the space $C_{1-\gamma}^{\alpha,\beta} [a, b]$ defined for $\gamma = \alpha + \beta - \alpha\beta$ ($0 < \alpha, \beta < 1$) by

$$C_{1-\gamma}^{\alpha,\beta} [a, b] = \left\{ y \in C_{1-\gamma} [a, b], D_{a+}^{\alpha,\beta} y \in C_{1-\gamma} [a, b] \right\}, \quad (3.2)$$

and

$$C_{1-\gamma}^{\gamma} [a, b] = \{ y \in C_{1-\gamma} [a, b], D_{a+}^{\gamma} y \in C_{1-\gamma} [a, b] \}.$$

We recall that $C_{1-\gamma} [a, b]$ is the weighted space of continuous functions on $(a, b]$

$$C_{1-\gamma} [a, b] = \left\{ g : (a, b] \rightarrow \mathbb{R} : (x - a)^{1-\gamma} g (x) \in C [a, b] \right\}, \quad (3.3)$$

and

$$\left(D_{a+}^{\alpha,\beta} y \right) (x) = \left(I_{a+}^{\beta(1-\alpha)} \frac{d}{dx} I_{a+}^{(1-\beta)(1-\alpha)} y \right) (x)$$

is the Hilfer fractional derivative of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$.

Our investigations are based on reducing the considered problem to a Volterra integral equation of the second kind

$$y(x) = \frac{c(x-a)^{(\alpha-1)(1-\beta)}}{\Gamma(\alpha+\beta-\alpha\beta)} + \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f[t, y(t)] dt, \quad x > a, \quad (3.4)$$

and on using the Banach fixed point theorem.

3.1 Equivalence of the Cauchy Type Problem and the Volterra Integral Equation

In this section we prove the equivalence between the Cauchy type problem (3.1) and the nonlinear Volterra integral equation (3.4) in the sense that, if $y \in C_{1-\gamma}^\gamma[a, b]$ satisfies one of these relations, then it also satisfies the other one. To establish such a result, we assume that a function $f[., y(.)]$ belongs to $C_{1-\gamma}[a, b]$ for any $y \in C_{1-\gamma}[a, b]$. For this we need the auxiliary assertion.

Lemma 3.1.1: *Let $\alpha > 0$, $0 \leq \gamma < 1$ and $f \in C_\gamma[a, b]$*

If $\gamma < \alpha$ then

$$(I_{a+}^\alpha f)(a) = \lim_{x \rightarrow a} (I_{a+}^\alpha f)(x) = 0, \quad 0 \leq \gamma < \alpha.$$

Proof : Since $f \in C_\gamma[a, b]$ then $(x-a)^\gamma f(x)$ is continuous on $[a, b]$ and on $[a, b]$ we

have

$$|(x - a)^\gamma f(x)| < M,$$

for some positive constant M .

Therefore

$$-M \left(I_{a+}^\alpha (t - a)^{-\gamma} \right) (x) < \left(I_{a+}^\alpha f \right) (x) < M \left(I_{a+}^\alpha (t - a)^{-\gamma} \right) (x)$$

and by using Property 2.2.1 (with $\beta = 1 - \gamma > 0$) we have

$$-M \frac{\Gamma(1-\gamma)}{\Gamma(\alpha+1-\gamma)} (x - a)^{\alpha-\gamma} < \left(I_{a+}^\alpha f \right) (x) < M \frac{\Gamma(1-\gamma)}{\Gamma(\alpha+1-\gamma)} (x - a)^{\alpha-\gamma} .$$

As $\alpha > \gamma$ we see that

$$\left(I_{a+}^\alpha f \right) (a) = \lim_{x \rightarrow a} \left(I_{a+}^\alpha f \right) (x) = 0, \quad 0 \leq \gamma < \alpha$$

which completes the proof of Lemma 3.1.1.

Theorem 3.1.1: Let $\gamma = \alpha + \beta - \alpha\beta$ where $0 < \alpha < 1$ and $0 \leq \beta \leq 1$. Let

$f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f[., y(.)] \in C_{1-\gamma}[a, b]$ for any $y \in C_{1-\gamma}[a, b]$.

If $y \in C_{1-\gamma}^\gamma[a, b]$, then y satisfies the (CFDP) (3.1) if and only if y satisfies the (IE) (3.4).

Proof : First we prove the **necessity**. Let $y \in C_{1-\gamma}^\gamma[a, b]$ be a solution of problem (3.1). We want to prove that y is also a solution of the integral equation (3.4). It is

clear, by the definition $C_{1-\gamma}^\gamma[a, b]$ above and Definition 2.2.3, that

$$D_{a+}^\gamma y = D(I_{a+}^{1-\gamma} y) \in C_{1-\gamma}[a, b], \quad I_{a+}^0 y = y,$$

and by Lemma 2.2.4(b) we have $I_{a+}^{1-\gamma} y \in C[a, b]$ since $y \in C_{1-\gamma}[a, b]$. Then by Definition 2.1.5 we have

$$I_{a+}^{1-\gamma} y \in C_{1-\gamma}^1[a, b].$$

Thus we can apply Lemma 2.2.8 to get

$$(I_{a+}^\gamma D_{a+}^\gamma y)(x) = y(x) - \frac{(I_{a+}^{1-\gamma} y)(a)}{\Gamma(\gamma)} (x-a)^{\gamma-1} \quad (3.5)$$

or

$$(I_{a+}^\gamma D_{a+}^\gamma y)(x) = y(x) - \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1}, \quad (3.6)$$

where c comes from the initial condition in (3.1). By our hypothesis $f[., y(.)] \in C_{1-\gamma}[a, b]$, and Lemma 2.2.1, we see that $I_{a+}^\alpha f \in C_{1-\gamma}[a, b]$. Applying the operator I_{a+}^α to both sides of (3.1) we get

$$I_{a+}^\alpha I_{a+}^{\beta(1-\alpha)} (D_{a+}^\gamma y) = I_{a+}^\alpha f[x, y(x)].$$

As we can apply lemma 2.2.6 to get

$$(I_{a+}^{\alpha+\beta(1-\alpha)} D_{a+}^\gamma y)(x) = I_{a+}^\alpha f[x, y(x)]$$

or

$$(I_{a+}^{\gamma} D_{a+}^{\gamma} y)(x) = (I_{a+}^{\alpha} f[t, y(t)])(x), \quad x \in (a, b]. \quad (3.7)$$

From (3.6) and (3.7) we obtain

$$y(x) = \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1} + (I_{a+}^{\alpha} f[t, y(t)])(x)$$

which is the equation (3.4), where $\gamma = \alpha + \beta - \alpha\beta$, and hence the necessity is proved.

Now we prove the **sufficiency**. Let $y \in C_{1-\gamma}^{\gamma}[a, b]$ satisfy the equation (3.4), then

$D_{a+}^{\gamma} y$ exists and $D_{a+}^{\gamma} y \in C_{1-\gamma}[a, b]$. Applying the operator D_{a+}^{γ} to both sides of (3.4)

we find

$$(D_{a+}^{\gamma} y)(x) = \frac{c}{\Gamma(\gamma)} (D_{a+}^{\gamma} (t-a)^{\gamma-1})(x) + (D_{a+}^{\gamma} I_{a+}^{\alpha} f[t, y(t)])(x).$$

By using Property 2.2.2 and Definition 2.2.3 when $0 < \gamma < 1$, we have

$$\begin{aligned} (D_{a+}^{\gamma} y)(x) &= \frac{d}{dx} (I_{a+}^{1-\gamma} I_{a+}^{\alpha} f[t, y(t)])(x) \\ &= \frac{d}{dx} \left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)] \right)(x) \\ &= \left(D_{a+}^{\beta(1-\alpha)} f[t, y(t)] \right)(x). \end{aligned} \quad (3.8)$$

From (3.8) and $D_{a+}^{\gamma} y \in C_{1-\gamma}[a, b]$, we obtain that

$$\left(D_{a+}^{\beta(1-\alpha)} f[t, y(t)] \right) \in C_{1-\gamma}[a, b].$$

Applying the operator $I_{a+}^{\beta(1-\alpha)}$ to both sides of (3.8) we get

$$\left(I_{a+}^{\beta(1-\alpha)} D_{a+}^{\gamma} y\right)(x) = \left(I_{a+}^{\beta(1-\alpha)} D_{a+}^{\beta(1-\alpha)} f[t, y(t)]\right)(x).$$

That is

$$I_{a+}^{\beta(1-\alpha)} \frac{d}{dx} \left(I_{a+}^{1-\gamma} y\right)(x) = \left(I_{a+}^{\beta(1-\alpha)} D_{a+}^{\beta(1-\alpha)} f[t, y(t)]\right)(x).$$

By virtue of $\frac{d}{dx} \left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)]\right) = \left(D_{a+}^{\beta(1-\alpha)} f[t, y(t)]\right) \in C_{1-\gamma}[a, b]$ and Lemma 2.2.4(b) we have $\left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)]\right) \in C[a, b]$ since $f[., y(.)] \in C_{1-\gamma}[a, b]$ with $1 - \gamma \leq 1 - \beta(1 - \alpha)$. Hence by Definition 2.1.5 we have

$$\left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)]\right) \in C_{1-\gamma}^1[a, b].$$

Then Lemma 2.2.8 allows us to write

$$\left(D_{a+}^{\alpha, \beta} y\right)(x) = f[x, y(x)] - \frac{\left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)]\right)(a)}{\Gamma[\beta(1-\alpha)]} (x-a)^{\beta(1-\alpha)-1}. \quad (3.9)$$

Lemma 3.1.1 (with γ replaced by $1-\gamma$, α by $1-\beta(1-\alpha)$ and $(1-\gamma) < 1-\beta(1-\alpha)$)

implies that

$$\left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)]\right)(a) = 0.$$

Hence the relation (3.9) reduces to

$$\left(D_{a+}^{\alpha, \beta} y\right) = f[x, y(x)].$$

Now we show that the initial condition in (3.1) also holds. To this end we apply the operator $I_{a+}^{1-\gamma}$ to both sides of (3.4)

$$(I_{a+}^{1-\gamma} y)(x) = \frac{c}{\Gamma(\gamma)} (I_{a+}^{1-\gamma} (t-a)^{\gamma-1})(x) + (I_{a+}^{1-\gamma} I_{a+}^{\alpha} f[t, y(t)])(x)$$

and use the Property 2.2.1 (with α replaced by $1-\gamma$ and β by γ) and Lemma 2.2.6 to obtain

$$(I_{a+}^{1-\gamma} y)(x) = c + \left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)] \right)(x). \quad (3.10)$$

In (3.10), taking the limit as $x \rightarrow a$, we obtain

$$(I_{a+}^{1-\gamma} y)(a) = c + \left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)] \right)(a).$$

But $\left(I_{a+}^{1-\beta(1-\alpha)} f[t, y(t)] \right)(a) = 0$ since $f[., y(.)] \in C_{1-\gamma}[a, b]$ (See Lemma 3.1.1) with γ replaced by $1-\gamma$ and α by $1-\beta(1-\alpha)$ where $1-\gamma < 1-\beta(1-\alpha)$. Therefore

$$(I_{a+}^{1-\gamma} y)(a) = c$$

and sufficiency is proved.

3.2 Existence and Uniqueness of the Global Solution to the Cauchy Type Problem

In this section we establish the existence of a unique solution to the Cauchy type problem (3.1) in the space $C_{1-\gamma}^{\alpha,\beta}[a, b]$ defined in (3.2) under the conditions of Theorem 3.1.1 and an additional Lipschitz condition on $f[., y(.)]$ with respect to the second variable: for all $x \in (a, b]$ and for all $y_1, y_2 \in \mathbf{G} \subset \mathbb{R}$,

$$|f[x, y_1] - f[x, y_2]| \leq A |y_1 - y_2| \quad (A > 0), \quad (3.11)$$

where $A > 0$ does not depend on $x \in (a, b]$.

We use the Banach fixed point theorem (see Theorem 2.5.1) to prove the existence and uniqueness in the appropriate space $C_{1-\gamma}^{\alpha,\beta}[a, b]$ where $\gamma = \alpha + \beta - \alpha\beta$.

Theorem 3.2.1: *Let $\gamma = \alpha + \beta - \alpha\beta$ where $(0 < \alpha < 1, 0 \leq \beta \leq 1)$. Let $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f[., y(.)] \in C_{1-\gamma}[a, b]$ for any $y \in C_{1-\gamma}[a, b]$ and the Lipschitzian condition (3.11) holds with respect to the second variable.*

Then there exists a unique solution y for the Cauchy type problem (3.1) in the space $C_{1-\gamma}^{\alpha,\beta}[a, b]$.

Proof : First we prove the existence of a unique solution y in the space $C_{1-\gamma}[a, b]$.

According to Theorem 3.1.1, it suffices to prove the existence of a unique solution $y \in C_{1-\gamma}[a, b]$ to the nonlinear Volterra integral equation (3.4). Let us select x_1 such

that the inequality

$$w_1 = \frac{A\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} (x_1 - a)^\alpha < 1, \quad (3.12)$$

where $A > 0$ is the Lipschitz constant in (3.11), is satisfied. We start by proving that a unique solution $y \in C_{1-\gamma}[a, x_1]$ to equation (3.4) exists on the interval $(a, x_1]$. For this, we use the Banach fixed point theorem for the space $C_{1-\gamma}[a, x_1]$, which is a complete metric space with the distance given by

$$d(y_1, y_2) = \|y_1 - y_2\|_{C_{1-\gamma}[a, x_1]} := \max_{x \in [a, x_1]} |(x - a)^{1-\gamma} [y_1(x) - y_2(x)]|.$$

The integral equation (3.4) takes the form

$$y(x) = (Ty)(x) \quad (3.13)$$

where

$$(Ty)(x) = y_0(x) + (I_{a+}^\alpha f[t, y(t)])(x) \quad (3.14)$$

with

$$y_0(x) = \frac{c}{\Gamma(\gamma)} (x - a)^{\gamma-1}. \quad (3.15)$$

Notice that T maps $C_{1-\gamma}[a, x_1]$ to itself. Since y_0 is clearly in $C_{1-\gamma}[a, x_1]$ and $(I_{a+}^\alpha f[t, y(t)])$ also is in $C_{1-\gamma}[a, x_1]$ by Lemma 2.2.1 when $y \in C_{1-\gamma}[a, x_1]$. Therefore $Ty \in C_{1-\gamma}[a, x_1]$.

Moreover T is a contraction, that is

$$\|Ty_1 - Ty_2\|_{C_{1-\gamma}[a, x_1]} \leq w_1 \|y_1 - y_2\|_{C_{1-\gamma}[a, x_1]}, \quad 0 < w_1 < 1. \quad (3.16)$$

This follows from (3.14), (3.11) and Lemma 2.2.1 as follows

$$\begin{aligned} \|Ty_1 - Ty_2\|_{C_{1-\gamma}[a, x_1]} &= \|I_{a+}^\alpha f[t, y_1(t)] - I_{a+}^\alpha f[t, y_2(t)]\|_{C_{1-\gamma}[a, x_1]} \\ &\leq \|I_{a+}^\alpha [f[t, y_1(t)] - f[t, y_2(t)]]\|_{C_{1-\gamma}[a, x_1]} \\ &\leq (x_1 - a)^\alpha \frac{\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} \|f[t, y_1(t)] - f[t, y_2(t)]\|_{C_{1-\gamma}[a, x_1]} \\ &\leq A(x_1 - a)^\alpha \frac{\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} \|y_1(t) - y_2(t)\|_{C_{1-\gamma}[a, x_1]} \\ &\leq w_1 \|y_1(t) - y_2(t)\|_{C_{1-\gamma}[a, x_1]}. \end{aligned}$$

Our assumption (3.12) allows us to apply the Banach fixed point Theorem (Theorem 2.5.1) to obtain a unique solution $y^* \in C_{1-\gamma}[a, x_1]$ to the equation (3.4) on the interval $[a, x_1]$.

Theorem 2.5.1 tells us that this solution y^* is a limit of a convergent sequence $T^m y_0^*$:

$$\lim_{m \rightarrow \infty} \|T^m y_0^* - y^*\|_{C_{1-\gamma}[a, x_1]} = 0,$$

where y_0^* is any function in $C_{1-\gamma}[a, x_1]$ and

$$(T^m y_0^*)(x) = (TT^{m-1} y_0^*)(x) = y_0(x) + (I_{a+}^\alpha f[t, (T^{m-1} y_0^*)(t)])(x), \quad m \in \mathbb{N}$$

It is convenient to take $y_0^*(x) = y_0(x)$ with $y_0(x)$ defined by (3.15). If we denote by

$$y_m(x) := (T^m y_0^*)(x), \quad m \in \mathbb{N}$$

then clearly

$$\lim_{m \rightarrow \infty} \|y_m - y^*\|_{C_{1-\gamma}[a, x_1]} = 0.$$

Next, we consider the interval $[x_1, b]$. From the equation (3.4) we have

$$\begin{aligned} y(x) &= \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1} + (I_{a+}^\alpha f[t, y(t)])(x) \\ &= \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f[t, y(t)] dt \\ &= \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^{x_1} (x-t)^{\alpha-1} f[t, y(t)] dt \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x (x-t)^{\alpha-1} f[t, y(t)] dt. \end{aligned}$$

Since the function $y(t)$ is uniquely defined on the interval $(a, x_1]$, the last integral can be considered as the unknown function, and we rewrite the last equation as

$$y(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x (x-t)^{\alpha-1} f[t, y(t)] dt, \quad (3.17)$$

where $y_{01}(x)$ is defined by

$$y_{01}(x) = \frac{c}{\Gamma(\gamma)} (x-a)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^{x_1} (x-t)^{\alpha-1} f[t, y^*(t)] dt, \quad (3.18)$$

and is a known function. We note that $y_{01}(x) \in C[x_1, b]$.

Next we prove the existence of a unique solution $y \in C[x_1, b]$ to equation (3.4) on

the interval $[x_1, b]$. For this, we also use Banach fixed point theorem for the space $C[x_1, x_2]$, where $x_2 \in (x_1, b]$ satisfies

$$w_2 = \frac{A}{\alpha \Gamma(\alpha)} (x_2 - x_1)^\alpha < 1. \quad (3.19)$$

$C[x_1, x_2]$ is a complete metric space with the distance given by

$$d(y_1, y_2) = \|y_1 - y_2\|_{C[x_1, x_2]} := \max_{x \in [x_1, x_2]} |y_1(x) - y_2(x)|.$$

The integral equation (3.17) may be written shortly as

$$y(x) = (Ty)(x), \quad (3.20)$$

where the operator (again denoted by T) is given by

$$(Ty)(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x (x-t)^{\alpha-1} f[t, y(t)] dt. \quad (3.21)$$

As in the first part of this proof, since $y_{01}(x) \in C[x_1, x_2]$ and $f[x, y(x)] \in C[x_1, x_2]$ for any $y \in C[x_1, x_2]$, then, by Lemma 2.2.2, We deduce that the integral in the right-hand side of (3.21) also belongs to $C[x_1, x_2]$, and hence $Ty \in C[x_1, x_2]$.

Moreover, using the Lipschitz condition (3.11) and applying Lemma 2.2.2, we find

$$\begin{aligned}
\|Ty_1 - Ty_2\|_{C[x_1, x_2]} &= \left\| I_{x_1^+}^\alpha f[t, y_1(t)] - I_{x_1^+}^\alpha f[t, y_2(t)] \right\|_{C[x_1, x_2]} \\
&\leq \left\| I_{x_1^+}^\alpha [|f[t, y_1(t)] - f[t, y_2(t)]|] \right\|_{C[x_1, x_2]} \\
&\leq \frac{1}{\alpha\Gamma(\alpha)} (x_2 - x_1)^\alpha \|f[t, y_1(t)] - f[t, y_2(t)]\|_{C[x_1, x_2]} \\
&\leq \frac{A}{\alpha\Gamma(\alpha)} (x_2 - x_1)^\alpha \|y_1(t) - y_2(t)\|_{C[x_1, x_2]} \\
&\leq w_2 \|y_1(t) - y_2(t)\|_{C[x_1, x_2]}.
\end{aligned}$$

This, together with our assumption $0 < w_2 < 1$, shows that T is a contraction and therefore from Theorem 2.5.1, there exists a unique solution $y_1^*(x) \in C[x_1, x_2]$ to equation (3.20) and hence to (3.4) on the interval $[x_1, x_2]$. Notice that $y_1^*(x_1) = y^*(x_1) = y_{01}(x_1)$. Furthermore Theorem 2.5.1, guarantees that this solution is the limit of a convergent sequence $T^m y_{01}^*$:

$$\lim_{m \rightarrow \infty} \|T^m y_{01}^* - y_1^*\|_{C[x_1, x_2]} = 0$$

where y_{01}^* is any function in $C[x_1, x_2]$, which we can pick $y_{01}^*(x) = y_{01}(x)$ defined by (3.18), if $y_0(x) \neq 0$ on $[x_1, x_2]$ we can take $y_{01}^*(x) = y_0(x)$ defined by (3.15).

Therefore

$$\lim_{m \rightarrow \infty} \|y_m - y_1^*\|_{C[x_1, x_2]} = 0,$$

where

$$y_m(x) = (T^m y_{01}^*)(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x (x-t)^{\alpha-1} f[t, (T^{m-1} y_{01}^*)(t)] dt.$$

If $x_2 \neq b$, we consider the interval $[x_2, x_3]$, such that $x_3 \leq b$ and

$$w_3 = \frac{A}{\alpha\Gamma(\alpha)} (x_3 - x_2)^\alpha < 1.$$

Using the same arguments as above, we derive that there exists a unique solution $y_2^*(x) \in C[x_2, x_3]$ to equation (3.4) on the interval $[x_2, x_3]$. If $x_3 \neq b$, then we continue the process until we reach a solution $y(x)$ to equation (3.4), $y(x) = y_k^*(x)$, and $y_k^*(x) \in C[x_k, x_{k+1}]$ ($k = 1, \dots, L$), where $a = x_0 < x_1 < \dots < x_{L+1}$ and

$$w_{k+1} = \frac{A}{\alpha\Gamma(\alpha)} (x_{k+1} - x_k)^\alpha < 1,$$

and we take $y_0(x) = y_{0k}(x)$, and $y_0^*(x) = y_{0k}^*(x)$ ($k = 1, \dots, L$) on each interval $[x_k, x_{k+1}]$. Assume that $b - a > \frac{\Gamma(\alpha+\gamma)}{A\Gamma(\gamma)}$ (for otherwise take $x_1 = b$). Then, divide the length of the interval $\left[a + \frac{\Gamma(\alpha+\gamma)}{A\Gamma(\gamma)}, b\right]$ by $\left(\frac{\alpha\Gamma(\alpha)}{A}\right)^{1/\alpha}$. Let M be that quotient. It is clear that $L = [M] + 1$ and b is reached after a finite number of steps, $x_{L+1} = b$. Then, there exists a unique solution $y(x) \in C[x_1, b]$ to equation (3.4) on the interval $[x_1, b]$.

Considering both parts of the solution in $[a, x_1]$ and $[x_1, b]$ as one single part on $[a, b]$ and taking into account Lemma 2.2.3, we obtain that there exists a unique solution $y(x) \in C_{1-\gamma}[a, b]$ to the Volterra integral equation (3.4) on the whole interval $[a, b]$, and hence $y \in C_{1-\gamma}[a, b]$ is the unique solution to the Cauchy-type problem (3.1).

It remains to show that such a unique solution $y \in C_{1-\gamma}[a, b]$ is actually in $C_{1-\gamma}^{\alpha, \beta}[a, b]$.

To this end we need to prove that $D_{a+}^{\alpha, \beta} y \in C_{1-\gamma}[a, b]$. Let us recall that our y is a

limit of the sequence $y_m(x)$, where $y_m(x) = (T^m y_0^*) \in C_{1-\gamma}[a, b]$ (where T is defined by (3.14) for $[a, x_1]$, by (3.21) for $[x_1, x_2]$, etc...), that is

$$\lim_{m \rightarrow \infty} \|y_m - y\|_{C_{1-\gamma}[a, b]} = 0,$$

with the choice of certain y_0^* on each $[a, x_1], \dots, [x_L, b]$. If $y_0 \neq 0$, then we can take $y_0^* = y_0$.

The equation in (3.1) and the Lipschitz condition (3.11), imply that

$$\begin{aligned} \left\| D_{a+}^{\alpha, \beta} y_m - D_{a+}^{\alpha, \beta} y \right\|_{C_{1-\gamma}[a, b]} &= \|f[x, y_m] - f[x, y]\|_{C_{1-\gamma}[a, b]} \\ &\leq A \|y_m - y\|_{C_{1-\gamma}[a, b]} \end{aligned}$$

and therefore

$$\lim_{m \rightarrow \infty} \left\| D_{a+}^{\alpha, \beta} y_m - D_{a+}^{\alpha, \beta} y \right\|_{C_{1-\gamma}[a, b]} = 0.$$

We entail from this relation that $D_{a+}^{\alpha, \beta} y \in C_{1-\gamma}[a, b]$ if $D_{a+}^{\alpha, \beta} y_m \in C_{1-\gamma}[a, b]$, $m = 1, 2, \dots$

This latter property holds from the relation

$$(T^m y_0^*)(x) = y_0(x) + (I_{a+}^{\alpha} f[t, (T^{m-1} y_0^*)(t)])(x), \quad m \in \mathbb{N}$$

and the fact that $f[x, y(x)] \in C_{1-\gamma}[a, b]$ for any $y \in C_{1-\gamma}[a, b]$. This completes the proof of Theorem 3.2.1.

3.3 Stability of Hilfer Fractional Derivative

In this section we are consider with the following fractional differential problem with weighted initial data

$$\begin{cases} D_{0+}^{\alpha,\beta} u(t) = f[t, u(t)], & t > 0, \\ t^{(1-\beta)(1-\alpha)} u(t) |_{t=0} = b, \end{cases} \quad (3.22)$$

where $D_{0+}^{\alpha,\beta}$ is the Hilfer fractional derivative (**HFD**) of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$, $b \in \mathbb{R}^*$ (the set of all real numbers except 0) and f is a continuous nonlinear function with respect to both of its arguments. We exploit the “initial decay” that is the behavior of solutions nearby the initial point and determine sufficient conditions on the nonlinearity which allow us to push and keep this behavior for all time in case of global existence.

We will assume the following hypotheses on the function $f(t, u)$:

(**F**) $f(t, u)$ is a continuous nonlinear function on $\mathbb{R}^+ \times \mathbb{R}$ and is such that

$$|f[t, u(t)]| \leq t^\mu e^{-\sigma t} \varphi(t) |u|^m, \quad \mu \geq 0, \quad m > 1, \quad \sigma > 0,$$

where φ is a continuous function on \mathbb{R}^+ . As $f[., y(.)]$ is continuous, under the assumption of Theorem 3.2.1 we have existence and uniqueness in the space $C_{1-\gamma}^{\alpha,\beta}[0, T]$ for any $T > 0$.

Let p and q be conjugate exponents, i.e. $pq = p + q$ and $\lambda_1 := 1 + p[\mu - (1 - \gamma)m]$,

$\lambda_2 := 1 + p(\alpha - 1)$ where $\gamma := \alpha + \beta - \alpha\beta$. If $\mu - (m - 1)(1 - \gamma) > 0$ and $q > 1/\alpha$, then $\lambda_1 > 0$ and $\lambda_2 > 0$. We denote by \mathbf{L} the positive real number

$$\mathbf{L} := \frac{1}{(m - 1) 2^{m(mq-1)} |b|^{mq(m-1)}} \left[\frac{\Gamma^p(\alpha) (\sigma p)^{\lambda_1}}{2^{p(1-\alpha)} \Gamma(\lambda_1) (1 + \lambda_1/\lambda_2)} \right]^{mq/p}.$$

Theorem 3.3.1: *Suppose that the hypotheses of Theorem 3.2.1 holds, $f[t, u(t)]$ satisfies **(F)** and $\mu - (m - 1)(1 - \gamma) > 0$. If $\varphi(t), \varphi(t)t^{-m\beta(1-\alpha)} \in L^q(0, \infty)$ for some $q > 1/\alpha$, then there exists a positive constant C such that the solution of (3.22) satisfies $|u(t)| \leq Ct^{\gamma-1}, t > 0$ where $\gamma = \alpha + \beta - \alpha\beta$, provided that*

$$\|\varphi\|_q^{(m-1)q} \left(\int_0^\infty s^{-qm\beta(1-\alpha)} \varphi^q(s) ds \right) < \mathbf{L}.$$

Proof : Let us consider the Volterra integral equation

$$u(t) = bt^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f[s, u(s)] ds, \quad t > 0, \quad (3.23)$$

associated to problem (3.22). Multiplying both sides of (3.23) by $t^{1-\gamma}$ and using the assumption **(F)** on $f([t, u(t)])$, we get

$$t^{1-\gamma} |u(t)| \leq |b| + \frac{t^{1-\gamma}}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} s^\mu e^{-\sigma s} \varphi(s) |u(s)|^m ds, \quad t > 0. \quad (3.24)$$

Let v denote the left-hand side of (3.24). Inserting the term $s^{(1-\gamma)m} s^{-(1-\gamma)m}$ inside

the integral gives

$$v(t) \leq |b| + \frac{t^{1-\gamma}}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} s^{\mu-(1-\gamma)m} e^{-\sigma s} \varphi(s) v^m(s) ds, \quad t > 0. \quad (3.25)$$

Now the Hölder inequality with exponents p and q yields

$$\begin{aligned} & \int_0^t (t-s)^{\alpha-1} s^{\mu-(1-\gamma)m} e^{-\sigma s} \varphi(s) v^m(s) ds \\ & \leq \left(\int_0^t (t-s)^{p(\alpha-1)} s^{p[\mu-(1-\gamma)m]} e^{-p\sigma s} ds \right)^{1/p} \left(\int_0^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > 0. \end{aligned}$$

The last term in the previous relation is finite for each t fixed. This follows from the fact φ is continuous and u is in $C_{1-\gamma}[0, T]$.

Since $\lambda_1 - 1 = p[\mu - (1 - \gamma)m]$, $\lambda_2 - 1 = p(\alpha - 1)$ and $\lambda_1, \lambda_2, p\sigma > 0$, we may apply Lemma 2.5.1 (with ν replaced by λ_2 , λ replaced by λ_1 and ω replaced by $p\sigma$) to get

$$\int_0^t (t-s)^{\alpha-1} s^{\mu-(1-\gamma)m} e^{-\sigma s} \varphi(s) v^m(s) ds \leq C_1 t^{\alpha-1} \left(\int_0^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad (3.26)$$

where C_1 is the constant appearing in Lemma 2.5.1 corresponding to the present exponents. That is

$$C_1 = \left[2^{p(1-\alpha)} \Gamma(\lambda_1) (1 + \lambda_1(\lambda_1 + 1)/\lambda_2) (p\sigma)^{-\lambda_1} \right]^{1/p}.$$

Combining (3.25) and (3.26) we entail that

$$v(t) \leq |b| + t^{-\beta(1-\alpha)} \hat{C}_1 \left(\int_0^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > 0, \quad (3.27)$$

where $\hat{C}_1 = C_1/\Gamma(\alpha)$. Multiplying both sides of (3.27) by $t^{\beta(1-\alpha)}$ we obtain

$$t^{\beta(1-\alpha)} v(t) \leq |b| t^{\beta(1-\alpha)} + \hat{C}_1 \left(\int_0^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > 0. \quad (3.28)$$

Let $z(t)$ denote the left-hand side of (3.28). Inserting the term $s^{-qm\beta(1-\alpha)} s^{qm\beta(1-\alpha)}$ inside the integral gives

$$z(t) \leq |b| t^{\beta(1-\alpha)} + \hat{C}_1 \left(\int_0^t \varphi^q(s) s^{-qm\beta(1-\alpha)} z^{qm}(s) ds \right)^{1/q}, \quad t > 0, \quad (3.29)$$

and raising both sides of (3.29) to the power q , we get (using Lemma 2.5.2)

$$z^q(t) \leq 2^{q-1} \left(|b|^q t^{q\beta(1-\alpha)} + \hat{C}_1^q \int_0^t \varphi^q(s) s^{-qm\beta(1-\alpha)} z^{qm}(s) ds \right), \quad t > 0. \quad (3.30)$$

Let us set

$$w(t) = \hat{C}_1^q \int_0^t \varphi^q(s) s^{-qm\beta(1-\alpha)} z^{qm}(s) ds, \quad t > 0. \quad (3.31)$$

Then, by the continuity of z and the assumption $\varphi(t) t^{-m\beta(1-\alpha)} \in L^q(0, \infty)$ the integrand is summable and $w(0) = 0$, and by differentiation

$$w'(t) = \hat{C}_1^q \varphi^q(t) t^{-qm\beta(1-\alpha)} z^{qm}(t). \quad (3.32)$$

Moreover, it is clear that φ and v are nonnegative continuous functions in \mathbb{R}^+ , and thus w is a continuous, nonnegative and nondecreasing function in \mathbb{R}^+ .

Now, we would like to estimate the right hand side of (3.32) in terms of w .

From (3.30) and (3.31) we entail that

$$z^q(t) \leq 2^{q-1} (|b|^q t^{q\beta(1-\alpha)} + w(t)), \quad t > 0.$$

Raising both sides to the power m and using Lemma 2.5.2, we get

$$z^{qm}(t) \leq 2^{mq-1} (|b|^{mq} t^{mq\beta(1-\alpha)} + w^m(t)). \quad (3.33)$$

Next, a substitution of (3.33) into (3.32) yields

$$\begin{aligned} w'(t) &\leq 2^{mq-1} \hat{C}_1^q \varphi^q(t) t^{-qm\beta(1-\alpha)} (|b|^{mq} t^{mq\beta(1-\alpha)} + w^m(t)) \\ &\leq 2^{mq-1} |b|^{mq} \hat{C}_1^q \varphi^q(t) + 2^{mq-1} \hat{C}_1^q t^{-qm\beta(1-\alpha)} \varphi^q(t) w^m(t), \quad t > 0. \end{aligned} \quad (3.34)$$

Applying Lemma 2.5.3 (with $w(u) = u^m$) we infer that

$$w(t) \leq G^{-1} [G(w(0) + l(t)) + k(t)],$$

where

$$l(t) = 2^{mq-1} |b|^{mq} \hat{C}_1^q \int_0^t \varphi^q(s) ds \text{ and } k(t) = 2^{mq-1} \hat{C}_1^q \int_0^t s^{-qm\beta(1-\alpha)} \varphi^q(s) ds. \text{ Since}$$

$G(r) = \int_{r_0}^r \frac{ds}{s^m}$, $r > 0$, $r_0 > 0$, then

$$G(r) = \frac{r^{1-m}}{1-m} - \frac{r_0^{1-m}}{1-m}$$

and

$$G^{-1}(y) = [r_0^{1-m} - (m-1)y]^{-\frac{1}{m-1}}.$$

That is

$$\begin{aligned} w(t) &\leq G^{-1} \left[\frac{l(t)^{1-m}}{1-m} - \frac{l(t_0)^{1-m}}{1-m} + k(t) \right] \\ &\leq \left[l(t_0)^{1-m} - (m-1) \left(\frac{l(t)^{1-m}}{1-m} - \frac{l(t_0)^{1-m}}{1-m} + k(t) \right) \right]^{-\frac{1}{m-1}} \\ &\leq [l(t)^{1-m} - (m-1)k(t)]^{-\frac{1}{m-1}} \end{aligned}$$

as long as

$$l(t)^{m-1} k(t) < \frac{1}{m-1}.$$

In particular, if $\left(\int_0^t \varphi^q(s) ds \right)^{m-1} \left(\int_0^t s^{-qm\beta(1-\alpha)} \varphi^q(s) ds \right) < \mathbf{L}$ then $w(t) \leq K_1$ for some positive constant K_1 , and thus from (3.29) we find that

$$z(t) \leq |b| t^{\beta(1-\alpha)} + K_1^{1/q}$$

or

$$t^{\beta(1-\alpha)} v(t) \leq |b| t^{\beta(1-\alpha)} + K_1^{1/q},$$

then

$$v(t) \leq |b| + K_1^{1/q} t^{-\beta(1-\alpha)} \leq C, \quad t \geq t_0 > 0$$

for some positive constant C . This yields that $|u(t)| \leq Ct^{\gamma-1}$ for $t \geq t_0 > 0$ and the proof is complete.

3.4 Non-existence of Solutions

In this section we consider the Cauchy problem of fractional order with a polynomial nonlinearity with variable coefficient

$$\begin{cases} \left(D_{0+}^{\alpha, \beta} u \right) (t) \geq t^\delta |u(t)|^m, & t > 0, m > 1, \delta \in \mathbb{R} \\ \left(D_{0+}^{\gamma-1} u \right) (0) = b > 0, \end{cases} \quad (3.35)$$

where $D_{0+}^{\alpha, \beta}$ is the Hilfer Fractional Derivative (**HFD**) of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$ and $\gamma = \alpha + \beta - \alpha\beta$.

There are several ways to investigate the non-existence and the blow up of solutions.

We will follow the paper by Laskri and Tatar [63], the proof is based on the test function method and the integration by parts formula.

For this we need the following lemma.

Lemma 3.4.1: *If $\alpha > 0$ and $f \in C[a, b]$, then*

$$\left(I_{a+}^\alpha f \right) (a) = \lim_{t \rightarrow a} \left(I_{a+}^\alpha f \right) (t) = 0$$

and

$$(I_{b-}^{\alpha} f)(b) = \lim_{t \rightarrow b} (I_{b-}^{\alpha} f)(t) = 0.$$

Proof : Since $f \in C[a, b]$, then on $[a, b]$, we have

$$|f(t)| < M,$$

for some positive constant M .

Therefore

$$\begin{aligned} |(I_{a+}^{\alpha} f)(t)| &\leq \frac{1}{\Gamma(\alpha)} \int_a^t (t-s)^{\alpha-1} |f(s)| ds \\ &\leq \frac{M}{\Gamma(\alpha)} \int_a^t (t-s)^{\alpha-1} ds \\ &\leq \frac{M}{\alpha \Gamma(\alpha)} [-(t-s)^{\alpha}]_{s=a}^t = \frac{M}{\alpha \Gamma(\alpha)} (t-a)^{\alpha}. \end{aligned}$$

As $\alpha > 0$ we see that

$$(I_{a+}^{\alpha} f)(a) = \lim_{t \rightarrow a} (I_{a+}^{\alpha} f)(t) = 0.$$

Similarity

$$\begin{aligned} |(I_{b-}^{\alpha} f)(t)| &\leq \frac{1}{\Gamma(\alpha)} \int_t^b (s-t)^{\alpha-1} |f(s)| ds \\ &\leq \frac{M}{\Gamma(\alpha)} \int_t^b (s-t)^{\alpha-1} ds \\ &\leq \frac{M}{\alpha \Gamma(\alpha)} [(s-t)^{\alpha}]_{s=t}^b = \frac{M}{\alpha \Gamma(\alpha)} (b-t)^{\alpha}. \end{aligned}$$

As $\alpha > 0$ we see that

$$(I_{b-}^{\alpha} f)(b) = \lim_{t \rightarrow b} (I_{b-}^{\alpha} f)(t) = 0.$$

Theorem (3.4.1): Assume that $\delta > -\alpha$ and $1 < m \leq \frac{\delta+1}{1-\alpha}$. Then, Problem (3.35)

does not admit global nontrivial solutions in $C_{1-\gamma}^\gamma[0, T]$, when $b > 0$.

Proof : Assume, on the contrary, that a nontrivial solution u exists for all time $t > 0$.

Let $\varphi \in C^1([0, \infty))$ be a test function satisfying : $\varphi(t) \geq 0$, φ is non-increasing such that

$$\varphi(t) := \begin{cases} 1, & t \in [0, T/2] \\ 0, & t \in [T, \infty) \end{cases}$$

for some $T > 0$. Multiplying the inequality in (3.35) by $\varphi(t)$ and integrating we get

$$\int_0^T \left(D_{0+}^{\alpha, \beta} u \right) (t) \varphi(t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt$$

and from the definition of $\left(D_{0+}^{\alpha, \beta} u \right) (t)$ we can write

$$\int_0^T I_{0+}^{\beta(1-\alpha)} \frac{d}{dt} \left(I_{0+}^{1-\gamma} u \right) (t) \varphi(t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt. \quad (3.36)$$

By virtue of Lemma 2.2.9, we may deduce from (3.36) that

$$\int_0^T \frac{d}{dt} \left(I_{0+}^{1-\gamma} u \right) (t) \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt. \quad (3.37)$$

An integration by parts in (3.37) yields

$$\left[\left(I_{0+}^{1-\gamma} u \right) (t) \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (t) \right]_{t=0}^T - \int_0^T \left(I_{0+}^{1-\gamma} u \right) (t) \frac{d}{dt} \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (t) dt$$

$$\geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt.$$

Note that the derivative inside the second term in the right hand side is meaningful because $\varphi \in [0, \infty)$ and $\varphi(T) = 0$ so the Riemann-Liouville and Caputo exist and are equal.

By using Lemma 3.4.1 we see that $\left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(T) = 0$ and also from initial condition $(I_{0+}^{1-\gamma}u)(0) = (D_{0+}^{\gamma-1}u)(0) = b$, so

$$-b \left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(0) - \int_0^T (I_{0+}^{1-\gamma}u)(t) \frac{d}{dt} \left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt, \quad (3.38)$$

It appears from Definition 2.2.4 that

$$-b \left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(0) + \int_0^T (I_{0+}^{1-\gamma}u)(t) \left(D_{T-}^{1-\beta(1-\alpha)}\varphi\right)(t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt \quad (3.39)$$

and from Lemma 2.2.5 we see that

$$\begin{aligned} & -b \left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(0) + \int_0^T (I_{0+}^{1-\gamma}u)(t) \left[\frac{1}{\Gamma[\beta(1-\alpha)]} \left(\frac{\varphi(T)}{(T-t)^{1-\beta(1-\alpha)}} \right. \right. \\ & \quad \left. \left. - \int_t^T \frac{\varphi'(s) ds}{(s-t)^{1-\beta(1-\alpha)}} \right) \right] \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt. \end{aligned} \quad (3.40)$$

Since $\varphi(T) = 0$ the relation (3.40) becomes

$$-b \left(I_{T-}^{\beta(1-\alpha)}\varphi\right)(0) - \int_0^T (I_{0+}^{1-\gamma}u)(t) \left(I_{T-}^{\beta(1-\alpha)}\varphi'\right)(t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt.$$

Lemma 2.2.9 again allows us to write

$$-b \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (0) - \int_0^T \varphi' (t) \left(I_{0+}^{\beta(1-\alpha)} I_{0+}^{1-\gamma} u \right) (t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt$$

and by Lemma 2.2.6

$$-b \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (0) - \int_0^T \varphi' (t) \left(I_{0+}^{1-\alpha} u \right) (t) dt \geq \int_0^T t^\delta |u(t)|^m \varphi(t) dt. \quad (3.41)$$

Notice that

$$\begin{aligned} \int_0^T \varphi' (t) \left(I_{0+}^{1-\alpha} u \right) (t) dt &= \frac{1}{\Gamma(1-\alpha)} \int_0^T \varphi' (t) \int_0^t \frac{u(s)}{(t-s)^\alpha} ds dt \\ &\leq \frac{1}{\Gamma(1-\alpha)} \int_0^T \left| \varphi' (t) \right| \int_0^t \frac{|u(s)|}{(t-s)^\alpha} ds dt. \end{aligned}$$

Since $\varphi(t)$ is nonincreasing, $\varphi(s) \geq \varphi(t)$ for all $t \geq s$, and thus

$$\frac{1}{\varphi(s)^{1/m}} \leq \frac{1}{\varphi(t)^{1/m}}, \quad 0 \leq s \leq t < T, \quad m > 1.$$

Also we have

$$\varphi' (t) = 0, \quad t \in [0, T/2].$$

Therefore

$$\begin{aligned}
\int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt &\leq \frac{1}{\Gamma(1-\alpha)} \int_0^T |\varphi'(t)| \int_0^t \frac{|u(s)|}{(t-s)^\alpha} \frac{\varphi(s)^{1/m}}{\varphi(s)^{1/m}} ds dt \\
&\leq \frac{1}{\Gamma(1-\alpha)} \int_0^T \frac{|\varphi'(t)|}{\varphi(t)^{1/m}} \int_0^t \frac{|u(s)|}{(t-s)^\alpha} \varphi(s)^{1/m} ds dt \\
&\leq \frac{1}{\Gamma(1-\alpha)} \int_{T/2}^T \frac{|\varphi'(t)|}{\varphi(t)^{1/m}} \int_0^t \frac{|u(s)|}{(t-s)^\alpha} \varphi(s)^{1/m} ds dt.
\end{aligned}$$

From Definition 2.2.1 we have

$$\int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt \leq \int_{T/2}^T \frac{|\varphi'(t)|}{\varphi(t)^{1/m}} (I_{0+}^{1-\alpha} [\varphi^{1/m} |u|])(t) dt$$

and by Lemma 2.2.9, we find

$$\int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt \leq \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right) (t) \varphi(t)^{1/m} |u(t)| dt.$$

(Note that we may assume that $|\varphi'(t)| \varphi(t)^{-1/m}$ is summable even though $\varphi(t) \rightarrow 0$

as $t \rightarrow T$, for otherwise we consider $\varphi^\lambda(t)$ with sufficiently large exponent λ).

Next, we multiply by $t^{\delta/m} \cdot t^{-\delta/m}$ inside the integral in the right hand side

$$\int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt \leq \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right) (t) \varphi(t)^{1/m} \frac{t^{\delta/m}}{t^{\delta/m}} |u(t)| dt.$$

For $-\alpha < \delta < 0$ we have $t^{-\delta/m} < T^{-\delta/m}$ (because $t < T$) and for $\delta > 0$ we get

$t^{-\delta/m} < 2^{\delta/m} T^{-\delta/m}$ (because $T/2 < t$), that is

$$t^{-\delta/m} < \max \{1, 2^{\delta/m}\} T^{-\delta/m}.$$

Therefore

$$\begin{aligned} & \int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt \\ & \leq \max \{1, 2^{\delta/m}\} T^{-\delta/m} \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right) (t) t^{\delta/m} \varphi(t)^{1/m} |u(t)| dt. \end{aligned} \quad (3.42)$$

A simple application of the Young inequality (Theorem 2.5.2) with m and m' such

that $\frac{1}{m} + \frac{1}{m'} = 1$ gives

$$\begin{aligned} & \int_0^T \varphi'(t) (I_{0+}^{1-\alpha} u)(t) dt \\ & \leq \frac{1}{m} \int_{T/2}^T t^\delta \varphi(t) |u(t)|^m dt + \frac{(\max \{1, 2^{\delta/m}\})^{m'}}{m'} T^{-\frac{\delta m'}{m}} \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right)^{m'} (t) dt \\ & \leq \frac{1}{m} \int_0^T t^\delta \varphi(t) |u(t)|^m dt + \frac{(\max \{1, 2^{\delta/m}\})^{m'}}{m'} T^{-\frac{\delta m'}{m}} \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right)^{m'} (t) dt. \end{aligned} \quad (3.43)$$

Clearly from (3.41) and (3.43) we see that

$$\begin{aligned} & -b \left(I_{T-}^{\beta(1-\alpha)} \varphi \right) (0) + \frac{(\max \{1, 2^{\delta/m}\})^{m'}}{m'} T^{-\frac{\delta m'}{m}} \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right)^{m'} (t) dt \\ & \geq \left(1 - \frac{1}{m} \right) \int_0^T t^\delta |u(t)|^m \varphi(t) dt, \end{aligned}$$

or since $b > 0$

$$\frac{1}{m'} \int_0^T t^\delta |u(t)|^m \varphi(t) dt \leq \frac{(\max \{1, 2^{\delta/m}\})^{m'}}{m'} T^{-\frac{\delta m'}{m}} \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right)^{m'} (t) dt. \quad (3.44)$$

Therefore, by Definition 2.2.2 we have

$$\begin{aligned} & \int_0^T t^\delta |u(t)|^m \varphi(t) dt \\ & \leq (\max\{1, 2^{\delta/m}\})^{m'} T^{-\frac{\delta m'}{m}} \int_{T/2}^T \left(\frac{1}{\Gamma(1-\alpha)} \int_t^T (s-t)^{-\alpha} \frac{|\varphi'(s)|}{\varphi(s)^{1/m}} ds \right)^{m'} dt. \end{aligned} \quad (3.45)$$

The change of variable $\sigma T = t$ yields

$$\begin{aligned} & \int_0^T t^\delta |u(t)|^m \varphi(t) dt \\ & \leq (\max\{1, 2^{\delta/m}\})^{m'} T^{-\frac{\delta m'}{m}} \int_{1/2}^1 \left(\frac{1}{\Gamma(1-\alpha)} \int_{\sigma T}^T (s-\sigma T)^{-\alpha} \frac{|\varphi'(s)|}{\varphi(s)^{1/m}} ds \right)^{m'} T d\sigma. \end{aligned} \quad (3.46)$$

Another change of variable $s = rT$ gives

$$\begin{aligned} & \int_0^T t^\delta |u(t)|^m \varphi(t) dt \\ & \leq (\max\{1, 2^{\delta/m}\})^{m'} T^{-\frac{\delta m'}{m}} \int_{1/2}^1 \left(\frac{1}{\Gamma(1-\alpha)} \int_{\sigma}^1 (rT - \sigma T)^{-\alpha} \frac{|\varphi'(r)|}{\varphi(r)^{1/m}} dr \right)^{m'} T d\sigma, \end{aligned}$$

or

$$\begin{aligned} & \int_0^T t^\delta |u(t)|^m \varphi(t) dt \\ & \leq \frac{(\max\{1, 2^{\delta/m}\})^{m'}}{\Gamma^{m'}(1-\alpha)} T^{1-\alpha m' - \delta m'/m} \int_{1/2}^1 \left(\int_{\sigma}^1 (r-\sigma)^{-\alpha} \frac{|\varphi'(r)|}{\varphi(r)^{1/m}} dr \right)^{m'} d\sigma. \end{aligned} \quad (3.47)$$

At this point it is clear that we may assume that the integral term in the right hand side of (3.47) is bounded, that is

$$\int_{1/2}^1 \left(\int_{\sigma}^1 (r - \sigma)^{-\alpha} \frac{|\varphi'(r)|}{\varphi(r)^{1/m}} dr \right)^{m'} d\sigma \leq K_1,$$

for some positive constant K_1 , for otherwise we consider $\varphi^\lambda(r)$ with some sufficiently large λ . Therefore

$$\int_0^T t^\delta |u(t)|^m \varphi(t) dt \leq K_2 T^{1-\alpha m' - \delta m'/m}, \quad (3.48)$$

with

$$K_2 := \frac{(\max\{1, 2^{\delta/m}\})^{m'}}{\Gamma^{m'}(1-\alpha)} K_1.$$

If $m < \frac{\delta+1}{1-\alpha}$ we see that $1 - \alpha m' - \delta m'/m < 0$ and consequently $T^{1-\alpha m' - \delta m'/m} \rightarrow 0$ as $T \rightarrow \infty$. Then from (3.48) we obtain

$$\lim_{T \rightarrow \infty} \int_0^T t^\delta |u(t)|^m \varphi(t) dt = 0.$$

We reach a contradiction since the solution is supposed to be nontrivial.

In the case $m = \frac{\delta+1}{1-\alpha}$ we have $1 - \alpha m' - \delta m'/m = 0$ and the relation (3.48) ensures that

$$\lim_{T \rightarrow \infty} \int_0^T t^\delta |u(t)|^m \varphi(t) dt \leq K_2. \quad (3.49)$$

Moreover, it is clear that

$$\begin{aligned} & \int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right) (t) t^{\delta/m} \varphi(t)^{1/m} |u(t)| dt \\ & \leq \left[\int_{T/2}^T \left(I_{T-}^{1-\alpha} \frac{|\varphi'|}{\varphi^{1/m}} \right)^{m'} (t) dt \right]^{\frac{1}{m'}} \left[\int_{T/2}^T t^{\delta} \varphi(t) |u(t)|^m dt \right]^{\frac{1}{m}}. \end{aligned}$$

This relation, together with (3.41) and (3.42), implies that

$$\int_0^T t^{\delta} \varphi(t) |u(t)|^m dt \leq K_3 \left[\int_{T/2}^T t^{\delta} \varphi(t) |u(t)|^m dt \right]^{\frac{1}{m}}$$

for some positive constant K_3 , with

$$\lim_{T \rightarrow \infty} \int_{T/2}^T t^{\delta} \varphi(t) |u(t)|^m dt = 0$$

due to the convergence of the integral in (3.49). This leads again to a contradiction.

This completes the proof of Theorem 3.4.1.

Chapter 4

Fractional Differential Problems with Hilfer-Hadamard Fractional Derivative

In this chapter we discuss the existence, uniqueness and solution, the stability of solutions of the Cauchy type problem (4.2).

Motivated by the definition of Hilfer fractional derivative which interpolates the Riemann-Liouville fractional derivative and the Caputo fractional derivative, we introduce next a similar one which interpolates the Hadamard fractional derivative (Definition 2.3.3) and its corresponding Hadamard-Caputo fractional derivative (Definition 2.4.2).

Definition (4.1): (*Hilfer–Hadamard Fractional Derivative (HHFD)*) *The left sided fractional derivative of order α ($0 < \alpha < 1$) and type $0 \leq \beta \leq 1$ with respect to x is defined by*

$$\left(\mathcal{D}_{a+}^{\alpha,\beta} f\right)(x) = \left(\mathcal{J}_{a+}^{\beta(1-\alpha)} \mathcal{D}_{a+}^{\alpha+\beta-\alpha\beta} f\right)(x) \quad (4.1)$$

where $\mathcal{D}_{a+}^{\alpha+\beta-\alpha\beta}$ is the Hadamard fractional derivative, for functions for which the expression on the right hand side exists.

Indeed for $\beta = 0$ this derivative (4.1) reduces to the Hadamard fractional derivative (Definition 2.3.3) and when $\beta = 1$ we recover the Hadamard-Caputo fractional derivative Definition (2.4.2).

We will study the existence and uniqueness of a solution of the Cauchy type problem

$$\begin{cases} \left(\mathcal{D}_{a+}^{\alpha,\beta} y\right)(x) = f(x, y), & x > a > 0 \\ \left(\mathcal{J}_{a+}^{(1-\beta)(1-\alpha)} y\right)(a) = c, \end{cases} \quad (4.2)$$

where $0 < \alpha < 1$, $0 \leq \beta \leq 1$ and $\mathcal{D}_{a^+}^{\alpha,\beta}$ is the **HHFD** (4.1) of order α and type β and c is a real number. We consider the underlying spaces $C_{\delta;1-\gamma,\mu}^{\alpha,\beta}[a, b]$ defined by

$$C_{\delta;1-\gamma,\mu}^{\alpha,\beta}[a, b] = \left\{ y \in C_{1-\gamma,\log}[a, b], \mathcal{D}_{a^+}^{\alpha,\beta} y \in C_{\mu,\log}[a, b] \right\} \quad (4.3)$$

and

$$C_{1-\gamma,\log}^\gamma[a, b] = \left\{ y \in C_{1-\gamma,\log}[a, b], \mathcal{D}_{a^+}^\gamma y \in C_{1-\gamma,\log}[a, b] \right\} \quad (4.4)$$

where $\gamma = \alpha + \beta - \alpha\beta$ and $0 \leq \mu < 1$. It is clear that $0 < \gamma < 1$ for $0 < \alpha, \beta < 1$.

Here $C_{1-\gamma,\log}[a, b]$ and $C_{\mu,\log}[a, b]$ are weighted spaces of continuous functions on $(a, b]$ defined by

$$C_{\gamma,\log}[a, b] = \left\{ g : (a, b] \rightarrow \mathbb{R} : \left(\log \frac{x}{a} \right)^\gamma g(x) \in C[a, b] \right\}, \quad (4.5)$$

Our investigations are based on reducing the fractional differential problem to a Volterra integral equation of the second kind

$$y(x) = \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a} \right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^x \left(\log \frac{x}{t} \right)^{\alpha-1} f[t, y(t)] \frac{dt}{t}, \quad x > a, \quad (4.6)$$

and on using the Banach fixed point theorem.

4.1 Equivalence of the Cauchy Type Problem and the Volterra Integral Equation

In this section we prove the equivalence of the Cauchy type problem (4.2) and the nonlinear Volterra integral equation (4.6) in the sense that, if $y \in C_{1-\gamma, \log}^\gamma[a, b]$ satisfies one of them, then it also satisfies the other one. To establish this result, we assume that the function $f[x, y(x)]$ belongs to $C_{\mu, \log}[a, b]$ for any $y \in \mathbf{G} \subset \mathbb{R}$. We will need the following lemma.

Lemma 4.1.1: *Let $0 < a < b < \infty$, $\alpha > 0$, $0 \leq \mu < 1$ and $g \in C_{\mu, \log}[a, b]$. If $\alpha > \mu$, then $\mathcal{J}_{a+}^\alpha g$ is continuous on $[a, b]$ and*

$$\mathcal{J}_{a+}^\alpha g(a) = \lim_{x \rightarrow a} \mathcal{J}_{a+}^\alpha g(x) = 0.$$

Proof : Since $g \in C_{\mu, \log}[a, b]$ then $(\log \frac{x}{a})^\mu g(x)$ is continuous on $[a, b]$ and on $[a, b]$ we have

$$\left| \left(\log \frac{x}{a} \right)^\mu g(x) \right| \leq M,$$

for some positive constant M . Therefore

$$|(\mathcal{J}_{a+}^\alpha g)(x)| \leq M \left(\mathcal{J}_{a+}^\alpha \left(\log \frac{x}{a} \right)^{-\mu} \right)(x)$$

and by using Property 2.3.1 (with $\beta = 1 - \mu > 0$) we have

$$|(\mathcal{J}_{a+}^\alpha g)(x)| \leq M \frac{\Gamma(1 - \mu)}{\Gamma(\alpha + 1 - \mu)} \left(\log \frac{x}{a} \right)^{\alpha - \mu}.$$

As $\alpha > \mu$, we obtain the result.

Theorem 4.1.1: *Let $\gamma = \alpha + \beta - \alpha\beta$ where $0 < \alpha < 1$ and $0 < \beta < 1$. Let $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ ($a > 0$) be a function such that $f(x, y) \in C_{\mu, \log}[a, b]$ for any $y \in C_{\mu, \log}[a, b]$ with $1 - \gamma \leq \mu < 1 - \beta(1 - \alpha)$.*

If $y \in C_{1-\gamma, \log}^\gamma[a, b]$, then y satisfies the (CFDP) (4.2) if and only if y satisfies the (IE) (4.6).

Proof : First we prove the **necessity**. Let $y \in C_{1-\gamma, \log}^\gamma[a, b]$ be a solution of problem (4.2). We want to prove that y is also a solution of the integral equation (4.6). By the definition of the space $C_{1-\gamma, \log}^\gamma[a, b]$ relation (4.4) above, we have

$$\delta(\mathcal{J}_{a^+}^{1-\gamma}y) = \mathcal{D}_{a^+}^\gamma y \in C_{1-\gamma, \log}[a, b].$$

Moreover, by Lemma 2.3.2(b) we have $\mathcal{J}_{a^+}^{1-\gamma}y \in C[a, b]$ since $y \in C_{1-\gamma, \log}[a, b]$. Then by Definition 2.1.7 we have

$$(\mathcal{J}_{a^+}^{1-\gamma}y)(x) \in C_{\delta, 1-\gamma}^1[a, b].$$

Thus we can apply Theorem 2.3.1 (with f replaced by y) to get

$$(\mathcal{J}_{a^+}^\gamma \mathcal{D}_{a^+}^\gamma y)(x) = y(x) - \frac{(\mathcal{J}_{a^+}^{1-\gamma}y)(a)}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1}, \quad x \in (a, b] \quad (4.7)$$

or

$$(\mathcal{J}_{a^+}^\gamma \mathcal{D}_{a^+}^\gamma y)(x) = y(x) - \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1}, \quad x \in (a, b] \quad (4.8)$$

where c comes from the initial condition in (4.2). By our hypothesis $f[x, y(x)] \in C_{\mu, \log}[a, b]$, since $y \in C_{1-\gamma, \log}[a, b] \subset C_{\mu, \log}[a, b]$, and Lemma 2.3.2 (a) and 2.3.2 (b) we see that the integral $\mathcal{J}_{a+}^{\alpha} f[x, y(x)] \in C_{\mu-\alpha, \log}[a, b]$ for $\mu > \alpha$ and $\mathcal{J}_{a+}^{\alpha} f[x, y(x)] \in C[a, b]$ for $\mu \leq \alpha$. Applying the operator $\mathcal{J}_{a+}^{\alpha}$ to both sides of (4.2) we get

$$\mathcal{J}_{a+}^{\alpha} \mathcal{J}_{a+}^{\beta(1-\alpha)} (\mathcal{D}_{a+}^{\gamma} y)(x) = \mathcal{J}_{a+}^{\alpha} f[x, y(x)], \quad x \in (a, b].$$

We can sum up the exponents by Property 2.3.2 to get

$$\mathcal{J}_{a+}^{\alpha+\beta(1-\alpha)} \mathcal{D}_{a+}^{\gamma} y(x) = \mathcal{J}_{a+}^{\alpha} f[x, y(x)], \quad x \in (a, b]$$

or

$$(\mathcal{J}_{a+}^{\gamma} \mathcal{D}_{a+}^{\gamma} y)(x) = (\mathcal{J}_{a+}^{\alpha} f[t, y(t)])(x), \quad x \in (a, b]. \quad (4.9)$$

From (4.8) and (4.9) we obtain

$$y(x) = \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a} \right)^{\gamma-1} + (\mathcal{J}_{a+}^{\alpha} f[t, y(t)])(x),$$

which is the equation (4.6), and hence the necessity is proved.

Now we prove the sufficiency. Let $y \in C_{1-\gamma, \log}^{\gamma}[a, b]$ satisfy the equation (4.6), then $\mathcal{D}_{a+}^{\gamma} y$ exists and $\mathcal{D}_{a+}^{\gamma} y \in C_{1-\gamma, \log}[a, b]$. Applying the operator $\mathcal{D}_{a+}^{\gamma}$ to both sides of (4.6) we find

$$(\mathcal{D}_{a+}^{\gamma} y)(x) = \frac{c}{\Gamma(\gamma)} \mathcal{D}_{a+}^{\gamma} \left(\log \frac{t}{a} \right)^{\gamma-1} (x) + (\mathcal{D}_{a+}^{\gamma} \mathcal{J}_{a+}^{\alpha} f[t, y(t)])(x).$$

By using Lemma 2.3.1, Definition 2.3.3, Property 2.3.2 and the hypothesis $f(x, y) \in C_{\mu, \log}[a, b]$, we have

$$\begin{aligned}
(\mathcal{D}_{a^+}^\gamma y)(x) &= \delta(\mathcal{J}_{a^+}^{1-\gamma} \mathcal{J}_{a^+}^\alpha f[t, y(t)])(x) \\
&= \delta(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f[t, y(t)])(x) \\
&= (\mathcal{D}_{a^+}^{\beta(1-\alpha)} f[t, y(t)])(x), \quad x \in (a, b].
\end{aligned} \tag{4.10}$$

From (4.10) and the fact that $\mathcal{D}_{a^+}^\gamma y \in C_{1-\gamma, \log}[a, b]$, we obtain that

$$\mathcal{D}_{a^+}^{\beta(1-\alpha)} f[x, y(x)] \in C_{1-\gamma, \log}[a, b].$$

Next, applying the operator $\mathcal{J}_{a^+}^{\beta(1-\alpha)}$ to both sides of (4.10) we get

$$(\mathcal{J}_{a^+}^{\beta(1-\alpha)} \mathcal{D}_{a^+}^\gamma y)(x) = (\mathcal{J}_{a^+}^{\beta(1-\alpha)} \mathcal{D}_{a^+}^{\beta(1-\alpha)} f[t, y(t)])(x).$$

That is

$$\mathcal{J}_{a^+}^{\beta(1-\alpha)} \delta(\mathcal{J}_{a^+}^{1-\gamma} y)(x) = (\mathcal{J}_{a^+}^{\beta(1-\alpha)} \mathcal{D}_{a^+}^{\beta(1-\alpha)} f[t, y(t)])(x).$$

By virtue of

$$\delta(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f[t, y(t)])(x) = (\mathcal{D}_{a^+}^{\beta(1-\alpha)} f[t, y(t)])(x) \in C_{1-\gamma, \log}[a, b],$$

and $\gamma > \beta(1-\alpha)$ and Definition 2.1.7 we have $\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f \in C_{\delta; 1-\gamma}^1[a, b]$ (see the first part of the proof for the continuity of $\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f$ for $\mu < 1 - \beta(1-\alpha)$). Then

Theorem 2.3.1 allows us to write

$$\left(\mathcal{D}_{a^+}^{\alpha,\beta} y\right)(x) = f(x, y) - \frac{\left(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f\right)(a)}{\Gamma[\beta(1-\alpha)]} \left(\log \frac{x}{a}\right)^{\beta(1-\alpha)-1}, \quad x \in (a, b]. \quad (4.11)$$

Lemma 4.1.1 implies that

$$\left(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f\right)(a) = 0$$

because $1 - \beta(1 - \alpha) > \mu$. Hence the relation (4.11) reduces to

$$\left(\mathcal{D}_{a^+}^{\alpha,\beta} y\right)(x) = f(x, y), \quad x > a.$$

Now we show that the initial condition in (4.2) also holds. To this end we apply the operator $\mathcal{J}_{a^+}^{1-\gamma}$ to both sides of (4.6)

$$\left(\mathcal{J}_{a^+}^{1-\gamma} y\right)(x) = \frac{c}{\Gamma(\gamma)} \mathcal{J}_{a^+}^{1-\gamma} \left(\log \frac{t}{a}\right)^{\gamma-1} (x) + \left(\mathcal{J}_{a^+}^{1-\gamma} \mathcal{J}_{a^+}^{\alpha} f[t, y(t)]\right)(x)$$

and use the Property 2.3.1 (with α replaced by $1 - \gamma$ and β by γ) and the Property 2.3.2 to obtain

$$\left(\mathcal{J}_{a^+}^{1-\gamma} y\right)(x) = c + \left(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f[t, y(t)]\right)(x). \quad (4.12)$$

In (4.12), taking the limit as $x \rightarrow a$, we obtain

$$\left(\mathcal{J}_{a^+}^{1-\gamma} y\right)(a) = c + \left(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f[t, y(t)]\right)(a).$$

But as mentioned above $\left(\mathcal{J}_{a^+}^{1-\beta(1-\alpha)} f[t, y(t)]\right)(a) = 0$, therefore

$$\left(\mathcal{J}_{a^+}^{1-\gamma} y\right)(a) = c$$

and the sufficiency is proved, which completes the proof of Theorem 4.1.1.

4.2 Existence and Uniqueness of a Solution

In this section we establish the existence of a unique solution to the Cauchy type problem (4.2) in the space $C_{\delta;1-\gamma,\mu}^{\alpha,\beta}[a, b]$ defined in (4.3) above under the conditions of Theorem 4.1.1 and an additional *Lipschitz* condition (Definition 2.5.1).

Theorem 4.2.1: *Let $\gamma = \alpha + \beta - \alpha\beta$ where $(0 < \alpha < 1, 0 \leq \beta \leq 1)$. Assume that $f : (a, b) \times \mathbb{R} \rightarrow \mathbb{R}$ ($a > 0$) is a function such that $f[x, y] \in C_{\mu,\log}[a, b]$ for any $y \in C_{\mu,\log}[a, b]$ with $1 - \gamma \leq \mu < 1 - \beta(1 - \alpha)$ and the Lipschitz condition (Definition 2.5.1) holds with respect to the second variable. Then there exists a unique solution y for the Cauchy type problem (4.2) in the space $C_{\delta;1-\gamma,\mu}^{\alpha,\beta}[a, b]$.*

Proof : First we prove the existence of a unique solution y in the space $C_{1-\gamma,\log}[a, b]$.

According to Theorem 4.1.1, it suffices to prove the existence of a unique solution $y \in C_{1-\gamma,\log}[a, b]$ to the nonlinear Volterra integral equation (4.6).

Let us select x_1 in (a, b) such that

$$w_1 := \frac{A\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} \left(\log \frac{x_1}{a}\right)^\alpha < 1, \quad (4.13)$$

where $A > 0$ is the Lipschitz constant in Definition 2.5.1. We start by proving that a unique solution $y \in C_{1-\gamma, \log} [a, x_1]$ to equation (4.6) exists on the interval $(a, x_1]$. It is easy to see that the space $C_{1-\gamma, \log} [a, x_1]$ is a complete metric space when equipped with the distance given by

$$d(y_1, y_2) = \|y_1 - y_2\|_{C_{1-\gamma, \log} [a, x_1]} := \max_{x \in [a, x_1]} \left| \left(\log \frac{x}{a} \right)^{1-\gamma} [y_1(x) - y_2(x)] \right|. \quad (4.14)$$

The integral equation (4.6) takes the form

$$y(x) = (Ty)(x) \quad (4.15)$$

where

$$(Ty)(x) = y_0(x) + (\mathcal{J}_{a^+}^\alpha f[t, y(t)])(x) \quad (4.16)$$

with

$$y_0(x) = \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a} \right)^{\gamma-1}. \quad (4.17)$$

We claim that T maps $C_{1-\gamma, \log} [a, x_1]$ to itself. Indeed, $y_0(x)$ given by (4.17) is clearly in $C_{1-\gamma, \log} [a, x_1]$. Also since $f[x, y(x)] \in C_{\mu, \log} [a, b]$ for any $y \in C_{\mu, \log} [a, b]$ with $\mu \in \mathbb{R}$ ($0 \leq \mu < 1$), then by Lemma 2.3.2 (a) and 2.3.2 (b) the integral in the right-hand side of (4.16) belongs to $C_{\mu-\alpha, \log} [a, b]$ for $\mu > \alpha$ and to $C[a, b]$ for $\mu \leq \alpha$. Since $\mu - \alpha < 1 - \gamma$, ($\mu < 1 - \beta(1 - \alpha)$), and $1 - \gamma \geq 0$. Then, by Property 2.1.1 the right-hand side of (4.16) belongs to $C_{1-\gamma, \log} [a, b]$. Therefore, the operator T defined by (4.16) maps $C_{1-\gamma, \log} [a, x_1]$ onto $C_{1-\gamma, \log} [a, x_1]$, that is $Ty \in C_{1-\gamma, \log} [a, x_1]$.

Our second claim is that T is a contraction, that is

$$\|Ty_1 - Ty_2\|_{C_{1-\gamma, \log}[a, x_1]} \leq w_1 \|y_1 - y_2\|_{C_{1-\gamma, \log}[a, x_1]}, \quad 0 < w_1 < 1. \quad (4.18)$$

This follows from (4.16), Definition 2.5.1, Lemma 2.3.2 (a) and the fact that

$$\begin{aligned} \|Ty_1 - Ty_2\|_{C_{1-\gamma, \log}[a, x_1]} &= \|\mathcal{J}_{a^+}^\alpha f[t, y_1(t)] - \mathcal{J}_{a^+}^\alpha f[t, y_2(t)]\|_{C_{1-\gamma, \log}[a, x_1]} \\ &\leq \|\mathcal{J}_{a^+}^\alpha [f[t, y_1(t)] - f[t, y_2(t)]]\|_{C_{1-\gamma, \log}[a, x_1]} \\ &\leq \left(\log \frac{x_1}{a}\right)^\alpha \frac{\Gamma(\gamma)}{\Gamma(\alpha+\gamma)} \|f[t, y_1(t)] - f[t, y_2(t)]\|_{C_{1-\gamma, \log}[a, x_1]} \\ &\leq A \left(\log \frac{x_1}{a}\right)^\alpha \frac{\Gamma(\gamma)}{\Gamma(\alpha+\gamma)} \|y_1(t) - y_2(t)\|_{C_{1-\gamma, \log}[a, x_1]} \\ &\leq w_1 \|y_1(t) - y_2(t)\|_{C_{1-\gamma, \log}[a, x_1]}. \end{aligned}$$

Our assumption (4.13) allows us to apply the Banach fixed point theorem (Theorem 2.5.1) to obtain a unique solution $y^* \in C_{1-\gamma, \log}[a, x_1]$ to the equation (4.6) on the interval $[a, x_1]$.

Theorem 2.5.1 tells us that this solution y^* is a limit of a convergent sequence $T^m y_0^*$:

$$\lim_{m \rightarrow \infty} \|T^m y_0^* - y^*\|_{C_{1-\gamma, \log}[a, x_1]} = 0, \quad (4.19)$$

where y_0^* is any function in $C_{1-\gamma, \log}[a, x_1]$. It is convenient to take $y_0^*(x) = y_0(x)$ with $y_0(x)$ defined by (4.17). If we denote by

$$y_m(x) := (T^m y_0^*)(x) = y_0(x) + \left(\mathcal{J}_{a^+}^\alpha f[t, (T^{m-1} y_0^*)(t)]\right)(x), \quad m \in \mathbb{N} \quad (4.20)$$

and $y_{m-1}(t) = (T^{m-1}y_0^*)(t)$, then clearly $\lim_{m \rightarrow \infty} \|y_m - y^*\|_{C_{1-\gamma, \log}[a, x_1]} = 0$.

Next, we consider the interval $[x_1, b]$. From the equation (4.6) we have

$$\begin{aligned} y(x) &= \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1} + \left(\mathcal{J}_{a+}^\alpha f[t, y(t)]\right)(x) \\ &= \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt \\ &= \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^{x_1} \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt \\ &\quad + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt. \end{aligned}$$

Since the function $y(t)$ is uniquely defined on the interval $(a, x_1]$, the last integral can be considered as the unknown function, and we rewrite the last equation as

$$y(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt, \quad (4.21)$$

where $y_{01}(x)$ is defined by

$$y_{01}(x) = \frac{c}{\Gamma(\gamma)} \left(\log \frac{x}{a}\right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^{x_1} \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt, \quad (4.22)$$

and is a known function. We note that $y_{01}(x) \in C[x_1, b]$.

Next, we consider the point $x_2 \in (x_1, b]$, $x_2 = x_1 + h_1$, $h_1 > 0$

$$w_2 = \frac{A}{\alpha \Gamma(\alpha)} \left(\log \frac{x_2}{x_1}\right)^\alpha < 1. \quad (4.23)$$

$C[x_1, x_2]$ is a complete metric space with the distance given by

$$d(y_1 - y_2) = \|y_1 - y_2\|_{C[x_1, x_2]} = \max_{x \in [x_1, x_2]} |y_1(x) - y_2(x)|. \quad (4.24)$$

The integral equation (4.21) may be written shortly as

$$y(x) = (Ty)(x), \quad (4.25)$$

where the operator (again denoted by T) is given by

$$(Ty)(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, y(t)]}{t} dt. \quad (4.26)$$

As in the first part of this proof, since $y_{01}(x) \in C[x_1, x_2]$ and $f[x, y(x)] \in C_{\mu, \log}[a, b]$ for any $y(x) \in C_{\mu, \log}[a, b]$, then $f[x, y(x)] \in C[x_1, x_2]$ for any $y(x) \in C[x_1, x_2]$, then by Lemma 2.3.4, we deduce that the integral in the right-hand side of (4.26) also belongs to $C[x_1, x_2]$, and hence $(Ty)(x) \in C[x_1, x_2]$.

Moreover, using the Lipschitz condition Definition 2.5.1 and applying the Lemma 2.3.4, we find

$$\begin{aligned} \|Ty_1 - Ty_2\|_{C[x_1, x_2]} &= \left\| \mathcal{J}_{x_1^+}^\alpha f[t, y_1(t)] - \mathcal{J}_{x_1^+}^\alpha f[t, y_2(t)] \right\|_{C[x_1, x_2]} \\ &\leq \left\| \mathcal{J}_{x_1^+}^\alpha |f[t, y_1(t)] - f[t, y_2(t)]| \right\|_{C[x_1, x_2]} \\ &\leq \frac{1}{\alpha \Gamma(\alpha)} \left(\log \frac{x_2}{x_1}\right)^\alpha \|f[t, y_1(t)] - f[t, y_2(t)]\|_{C[x_1, x_2]} \\ &\leq \frac{A}{\alpha \Gamma(\alpha)} \left(\log \frac{x_2}{x_1}\right)^\alpha \|y_1(t) - y_2(t)\|_{C[x_1, x_2]} \\ &\leq w_2 \|y_1(t) - y_2(t)\|_{C[x_1, x_2]}. \end{aligned} \quad (4.27)$$

This, together with our assumption $0 < w_2 < 1$, shows that T is a contraction and therefore from Theorem 2.5.1, there exists a unique solution $y_1^*(x) \in C[x_1, x_2]$ to equation (4.6) on the interval $[x_1, x_2]$. Notice that $y_1^*(x_1) = y^*(x_1) = y_{01}(x_1)$. Further, Theorem 2.5.1 guarantees that this solution is the limit of a convergent sequence $T^m y_{01}^*$:

$$\lim_{m \rightarrow \infty} \|T^m y_{01}^* - y_1^*\|_{C[x_1, x_2]} = 0 \quad (4.28)$$

where y_{01}^* is any function in $C[x_1, x_2]$, which we can pick $y_{01}^*(x) = y_{01}(x)$ defined by (4.22). Therefore

$$\lim_{m \rightarrow \infty} \|y_m - y_1^*\|_{C[x_1, x_2]} = 0, \quad (4.29)$$

where

$$y_m(x) = (T^m y_{01}^*)(x) = y_{01}(x) + \frac{1}{\Gamma(\alpha)} \int_{x_1}^x \left(\log \frac{x}{t}\right)^{\alpha-1} \frac{f[t, (T^{m-1} y_{01}^*)(t)]}{t} dt. \quad (4.30)$$

If $x_2 \neq b$, we consider the interval $[x_2, x_3]$, where $x_3 = x_2 + h_2$, $h_2 > 0$ such that $x_3 \leq b$ and

$$w_3 = \frac{A}{\alpha \Gamma(\alpha)} \left(\log \frac{x_3}{x_2}\right)^\alpha < 1. \quad (4.31)$$

Using the same arguments as above, we derive that there exists a unique solution $y_2^* \in C[x_2, x_3]$ to the equation (4.6) on the interval $[x_2, x_3]$. If $x_3 \neq b$, then we continue the process until we reach a solution y to (4.6), $y(x) = y_k^*(x)$, and $y_k^* \in C[x_k, x_{k+1}]$ ($k = 1, \dots, L$), where $a = x_0 < x_1 < \dots < x_{L+1}$ and

$$w_{k+1} = \frac{A}{\alpha \Gamma(\alpha)} \left(\log \frac{x_{k+1}}{x_k}\right)^\alpha < 1, \quad (4.32)$$

Assume that $b - a > \frac{\Gamma(\alpha+\gamma)}{A\Gamma(\gamma)}$ (for otherwise take $x_1 = b$). Then, divide the length of the interval $\left[a + \frac{\Gamma(\alpha+\gamma)}{A\Gamma(\gamma)}, b\right]$ by $\left(\frac{\alpha\Gamma(\alpha)}{A}\right)^{1/\alpha}$. Let M be that quotient. It appears that for $L = [M] + 1$ and b is reached after a finite number of steps, $x_{L+1} = b$. Then, there exists a unique solution $y(x) \in C[x_1, b]$ to equation (4.6) on the interval $[x_1, b]$.

Putting together the solutions in $[a, x_1]$ and $[x_1, b]$ and taking into account the Lemma 2.3.5, we obtain that there exists a unique solution $y \in C_{1-\gamma, \log}[a, b]$ to the Volterra integral equation (4.6) on the whole interval $[a, b]$. Hence $y \in C_{1-\gamma, \log}[a, b]$ is the unique solution to the Cauchy-type problem (4.2).

It remains to show that such a unique solution $y \in C_{1-\gamma, \log}[a, b]$ is actually in $C_{\delta; 1-\gamma, \mu}^{\alpha, \beta}[a, b]$. To this end we need to prove that $\mathcal{D}_{a^+}^{\alpha, \beta} y \in C_{\mu, \log}[a, b]$. Let us recall that our y is a limit of the sequence y_m , where $y_m(x) = (T^m y_0^*) \in C_{1-\gamma, \log}[a, b]$, that is

$$\lim_{m \rightarrow \infty} \|y_m - y\|_{C_{1-\gamma, \log}[a, b]} = 0, \quad (4.33)$$

with a certain choice of $y_0^*(x)$ on each $[a, x_1], \dots, [x_L, b]$.

If $y_0(x) \neq 0$, then we can take $y_0^*(x) = y_0(x)$. Since $\mu \geq 1 - \gamma$, then by (4.2), the Lipschitz condition Definition 2.5.1 and Property 2.1.1, we have

$$\begin{aligned} \left\| \mathcal{D}_{a^+}^{\alpha, \beta} y_m - \mathcal{D}_{a^+}^{\alpha, \beta} y \right\|_{C_{\mu, \log}[a, b]} &= \|f[x, y_m] - f[x, y]\|_{C_{\mu, \log}[a, b]} \\ &\leq A \|y_m - y\|_{C_{\mu, \log}[a, b]} \leq A \left(\log \frac{b}{a}\right)^{\mu-1+\gamma} \|y_m - y\|_{C_{1-\gamma, \log}[a, b]}. \end{aligned} \quad (4.34)$$

In virtue of (4.33) and (4.34) it follows that

$$\lim_{m \rightarrow \infty} \left\| \mathcal{D}_{a^+}^{\alpha, \beta} y_m - \mathcal{D}_{a^+}^{\alpha, \beta} y \right\|_{C_{\mu, \log}[a, b]} = 0. \quad (4.35)$$

We entail from this relation that $\left(\mathcal{D}_{a^+}^{\alpha,\beta} y\right)(x) \in C_{\mu,\log}[a, b]$ if $\left(\mathcal{D}_{a^+}^{\alpha,\beta} y_m\right)(x) \in C_{\mu,\log}[a, b]$, $m = 1, 2, \dots$. This latter property holds in as much as $\mathcal{D}_{a^+}^{\alpha,\beta} y_m(x) = f[x, y_{m-1}(x)]$ and $f[x, y(x)] \in C_{\mu,\log}[a, b]$ for any $y(x) \in C_{\mu,\log}[a, b]$. Hence a unique solution $y(x) \in C_{1-\gamma,\log}[a, b]$ has the property $\left(\mathcal{D}_{a^+}^{\alpha,\beta} y\right)(x) \in C_{\mu,\log}[a, b]$. Consequently, $y(x) \in C_{\delta;1-\gamma,\mu}^{\alpha,\beta}[a, b]$. This completes the proof of Theorem 4.2.1.

4.3 Stability of Hilfer-Hadamard Fractional Derivative

In this section we consider the weighted Cauchy-type problem

$$\begin{cases} \mathcal{D}_{a^+}^{\alpha,\beta} u(t) = f(t, u), & t > a > 0, \\ \left(\log \frac{t}{a}\right)^{(1-\beta)(1-\alpha)} u(t) \big|_{t=a} = b, \end{cases} \quad (4.36)$$

where $\mathcal{D}_+^{\alpha,\beta}$ is the Hilfer-Hadamard fractional derivative (**HHFD**) of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$ and $b \in \mathbb{R}^*$ (the set of all real numbers except 0). We will assume the following hypotheses on the function $f(t, u)$:

(**F***) $f(t, u)$ is a continuous nonlinear function on $(a, \infty) \times \mathbb{R}$ and is such that

$$|f(t, u)| \leq \left(\log \frac{t}{a}\right)^\mu \varphi(t) |u(t)|^m, \quad \mu \geq 0, \quad m > 1, \quad t \geq a$$

where φ is a continuous function on $[a, \infty)$.

For this we need the following inequality.

Lemma 4.3.1 : *If $\lambda, \nu, \omega > 0$, then for any $t > a, a > 0$ we have*

$$\left(\log \frac{t}{a}\right)^{1-\nu} \int_a^t \left(\log \frac{t}{s}\right)^{\nu-1} \left(\log \frac{s}{a}\right)^{\lambda-1} \left(\frac{s}{a}\right)^{-\omega} \frac{ds}{s} \leq C\omega^{-\lambda},$$

where C is a positive constant independent of t .

Proof : Let us denote by $I(t)$ the left-hand side of the inequality in the Lemma. We

consider the change of variable, $\xi = \frac{\log(s/a)}{\log(t/a)}$ then $\frac{s}{a} = \left(\frac{t}{a}\right)^\xi$ and $\log \frac{t}{s} = (1 - \xi) \left(\log \frac{t}{a}\right)$.

It follows that

$$I(t) = \left(\log \frac{t}{a}\right)^\lambda \int_0^1 (1 - \xi)^{\nu-1} \xi^{\lambda-1} \left(\frac{t}{a}\right)^{-\omega\xi} d\xi$$

or

$$I(t) = \left(\log \frac{t}{a}\right)^\lambda \int_0^1 (1 - \xi)^{\nu-1} \xi^{\lambda-1} \exp\left(-\omega\xi \log \left(\frac{t}{a}\right)\right) d\xi.$$

Observe that, for $\xi \geq 1$ and $[\lambda] + 1 \geq \lambda$ we have $\xi^{[\lambda]+1} \geq \xi^\lambda$. Also since $\lambda + 2 \geq [\lambda] + 2$

and the Gamma function is increasing for $[2, \infty)$, we have $\Gamma(\lambda + 2) \geq \Gamma([\lambda] + 2)$ or

$\frac{1}{\Gamma([\lambda]+2)} \geq \frac{1}{\Gamma(\lambda+2)}$. Moreover $e^\xi \geq \frac{\xi^{[\lambda]+1}}{\Gamma([\lambda]+2)}$ ($\Gamma([\lambda] + 2) = ([\lambda] + 1)!$), and hence

$$e^\xi \geq \frac{\xi^{[\lambda]+1}}{\Gamma([\lambda] + 2)} \geq \frac{\xi^\lambda}{\Gamma([\lambda] + 2)} \geq \frac{\xi^\lambda}{\Gamma(\lambda + 2)}$$

or

$$e^{-\xi} \leq \frac{\Gamma(\lambda + 2)}{\xi^\lambda}.$$

Therefore, for $0 \leq \xi < 1/2$ we get

$$(1 - \xi)^{\nu-1} \leq \max(1, 2^{1-\nu}).$$

For $1/2 < \xi \leq 1$ we have

$$\exp\left(-\omega\xi \log\left(\frac{t}{a}\right)\right) \leq \frac{\Gamma(\lambda+2)}{(\omega\xi \log(\frac{t}{a}))^\lambda} \leq \frac{\omega^{-\lambda}}{\xi} \Gamma(\lambda+2) \leq 2\omega^{-\lambda} \Gamma(\lambda+2).$$

This means that

$$\begin{aligned} & \left(\log \frac{t}{a}\right)^\lambda (1 - \xi)^{\nu-1} \xi^{\lambda-1} \exp\left(-\omega\xi \log\left(\frac{t}{a}\right)\right) \\ & \leq \begin{cases} \max(1, 2^{1-\nu}) \left(\log \frac{t}{a}\right)^\lambda \xi^{\lambda-1} \exp\left(-\omega\xi \log\left(\frac{t}{a}\right)\right) & \text{for } 0 \leq \xi < 1/2, \\ 2(1 - \xi)^{\nu-1} \Gamma(\lambda+2) \omega^{-\lambda} & \text{for } 1/2 < \xi \leq 1. \end{cases} \end{aligned}$$

Consequently

$$\begin{aligned} I(t) & \leq \max(1, 2^{1-\nu}) \left(\log \frac{t}{a}\right)^\lambda \int_0^{1/2} \xi^{\lambda-1} \exp\left(-\omega\xi \log\left(\frac{t}{a}\right)\right) d\xi \\ & \quad + 2\omega^{-\lambda} \Gamma(\lambda+2) \int_{1/2}^1 (1 - \xi)^{\nu-1} d\xi. \end{aligned}$$

Let $u = \omega\xi \log\left(\frac{t}{a}\right)$, we see that

$$I(t) \leq \max(1, 2^{1-\nu}) \left(\log \frac{t}{a}\right)^\lambda \int_0^\infty \left(\frac{u}{\omega \log\left(\frac{t}{a}\right)}\right)^{\lambda-1} e^{-u} \frac{du}{\omega \log\left(\frac{t}{a}\right)}$$

$$+2\omega^{-\lambda}\Gamma(\lambda+2)\left[-\frac{(1-\xi)^\nu}{\nu}\right]_{\xi=1/2}^1.$$

Thus

$$I(t) \leq \max(1, 2^{1-\nu}) \omega^{-\lambda} \Gamma(\lambda) + \frac{2^{1-\nu} \omega^{-\lambda} \Gamma(\lambda+2)}{\nu}.$$

As a result, $I(t) \leq \max\{1, 2^{1-\nu}\} \Gamma(\lambda) (1 + \lambda(\lambda+1)/\nu) \omega^{-\lambda}$.

For $0 < \xi < 1$, $e^\xi \geq 1$ it is clear that

$$\Gamma(\lambda+2) e^\xi \geq 1 \geq \xi^\lambda$$

holds and we proceed in the same manner to conclude that

$$\left(\log \frac{t}{a}\right)^{1-\nu} \int_a^t \left(\log \frac{t}{s}\right)^{\nu-1} \left(\log \frac{s}{a}\right)^{\lambda-1} \left(\frac{s}{a}\right)^{-\omega} \frac{ds}{s} \leq C \omega^{-\lambda},$$

where $C = \max\{1, 2^{1-\nu}\} \Gamma(\lambda) (1 + \lambda(\lambda+1)/\nu)$. The proof is complete.

Let p and q be conjugate exponents, i.e. $pq = p+q$, and let $\lambda_1 := 1+p[\mu - (1-\gamma)m]$

and $\lambda_2 := 1+p(\alpha-1)$, where $\gamma := \alpha+\beta-\alpha\beta$. If $\mu-(m-1)(1-\gamma) > 0$ and $q > 1/\alpha$,

then $\lambda_1 > 0$ and $\lambda_2 > 0$. We denote by \mathbf{L}^* the positive real number

$$\mathbf{L}^* := \left(\frac{\Gamma(\alpha)}{2^{m+(\alpha-1)}|b|^{m-1}}\right)^m \left(\frac{(2a)^m}{m-1}\right)^{1/q} \left[\frac{(p-1)^{\lambda_1}}{\Gamma(\lambda_1)(1+\lambda_1/\lambda_2)}\right]^{m/p}, \quad m > 1.$$

Theorem 4.3.1: Suppose that $f(t, u)$ satisfies (\mathbf{F}^*) and $\mu > (m - 1)(1 - \gamma)$. If

$$\left(\|\varphi(t)\|_q \right)^{m-1} \left\| \varphi(t) \left(\log \frac{t}{a} \right)^{-m\beta(1-\alpha)} \right\|_q < \mathbf{L}^*$$

for some $q > 1/\alpha$, then there exists a positive constant C such that $|u(t)| \leq C \left(\log \frac{t}{a} \right)^{\gamma-1}$, $t > ae$ ($a > 0$), where $\gamma = \alpha + \beta - \alpha\beta$.

Proof : Let us consider the Volterra integral equation

$$u(t) = b \left(\log \frac{t}{a} \right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} f[s, u(s)] \frac{ds}{s}, \quad t > a > 0 \quad (4.37)$$

associated to problem (4.36). Multiplying both sides of (4.37) by $\left(\log \frac{t}{a} \right)^{1-\gamma}$ and using the assumption (\mathbf{F}^*) on $f(t, u)$, we get

$$\left(\log \frac{t}{a} \right)^{1-\gamma} |u(t)| \leq |b| + \frac{\left(\log \frac{t}{a} \right)^{1-\gamma}}{\Gamma(\alpha)} \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \left(\log \frac{s}{a} \right)^\mu \varphi(s) |u(s)|^m \frac{ds}{s}. \quad (4.38)$$

Let $v(t)$ denote the left-hand side of (4.38). The insertion of the term

$$\left(\log \frac{s}{a} \right)^{(1-\gamma)m} \left(\log \frac{s}{a} \right)^{-(1-\gamma)m}$$

inside the integral gives

$$v(t) \leq |b| + \frac{\left(\log \frac{t}{a} \right)^{1-\gamma}}{\Gamma(\alpha)} \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \left(\log \frac{s}{a} \right)^{\mu-(1-\gamma)m} \varphi(s) v^m(s) \frac{ds}{s}, \quad t > a. \quad (4.39)$$

Now the Hölder inequality with exponents p and q yields

$$\begin{aligned} & \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \left(\log \frac{s}{a} \right)^{\mu-(1-\gamma)m} \varphi(s) v^m(s) \frac{ds}{s} \\ & \leq \left(\int_a^t \left(\log \frac{t}{s} \right)^{p(\alpha-1)} \left(\log \frac{s}{a} \right)^{p(\mu-(1-\gamma)m)} \frac{ds}{s^p} \right)^{1/p} \left(\int_a^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \end{aligned}$$

or

$$\begin{aligned} & \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \left(\log \frac{s}{a} \right)^{\mu-(1-\gamma)m} \varphi(s) v^m(s) \frac{ds}{s} \\ & \leq a^{-1/q} \left(\int_a^t \left(\log \frac{t}{s} \right)^{p(\alpha-1)} \left(\log \frac{s}{a} \right)^{p(\mu-(1-\gamma)m)} \left(\frac{s}{a} \right)^{-(p-1)} \frac{ds}{s} \right)^{1/p} \\ & \quad \times \left(\int_a^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > a. \end{aligned}$$

Since $\lambda_1 - 1 = p[\mu - (1 - \gamma)m]$, $\lambda_2 - 1 = p(\alpha - 1)$ and $\lambda_1, \lambda_2, p - 1 > 0$, we may apply Lemma 4.3.1 (with ν replaced by λ_2 , λ replaced by λ_1 and ω replaced by $p - 1$) to get

$$\begin{aligned} & \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \left(\log \frac{s}{a} \right)^{\mu-(1-\gamma)m} \varphi(s) v^m(s) \frac{ds}{s} \leq a^{-1/q} C_1 \left(\log \frac{t}{a} \right)^{\alpha-1} \\ & \quad \times \left(\int_a^t \varphi^q(s) v^{qm}(s) ds \right)^{\frac{1}{q}}, \quad t > a \end{aligned} \tag{4.40}$$

where C_1 is the constant appearing in Lemma 4.3.1 corresponding to the present exponents. That is

$$C_1 = \left(2^{p(\alpha-1)} \Gamma(\lambda_1) (1 + \lambda_1(\lambda_1 + 1)/\lambda_2) (p - 1)^{-\lambda_1} \right)^{1/p}.$$

Combining (4.39) and (4.40) we entail that

$$v(t) \leq |b| + \hat{C}_1 \left(\log \frac{t}{a} \right)^{-\beta(1-\alpha)} \left(\int_a^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > a \quad (4.41)$$

where $\hat{C}_1 = a^{-1/q} \frac{C_1}{\Gamma(\alpha)}$. Multiplying both sides of (4.41) by $\left(\log \frac{t}{a} \right)^{\beta(1-\alpha)}$, we obtain

$$\left(\log \frac{t}{a} \right)^{\beta(1-\alpha)} v(t) \leq |b| \left(\log \frac{t}{a} \right)^{\beta(1-\alpha)} + \hat{C}_1 \left(\int_a^t \varphi^q(s) v^{qm}(s) ds \right)^{1/q}, \quad t > a. \quad (4.42)$$

Let $z(t)$ denote the left-hand side of (4.42). The insertion of the term

$$\left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} \left(\log \frac{s}{a} \right)^{qm\beta(1-\alpha)}$$

inside the integral gives

$$z(t) \leq |b| \left(\log \frac{t}{a} \right)^{\beta(1-\alpha)} + \hat{C}_1 \left(\int_a^t \varphi^q(s) \left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} z^{qm}(s) ds \right)^{1/q}. \quad (4.43)$$

Raising both sides of (4.43) to the power q and using Lemma 2.5.2, we get

$$z^q(t) \leq 2^{q-1} \left(|b|^q \left(\log \frac{t}{a} \right)^{q\beta(1-\alpha)} + \hat{C}_1^q \int_a^t \varphi^q(s) \left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} z^{qm}(s) ds \right). \quad (4.44)$$

Let us set

$$w(t) = \hat{C}_1^q \int_a^t \varphi^q(s) \left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} z^{qm}(s) ds, \quad t > a. \quad (4.45)$$

Then, clearly $w(a) = 0$, and by differentiation

$$w'(t) = \hat{C}_1^q \varphi^q(t) \left(\log \frac{t}{a} \right)^{-qm\beta(1-\alpha)} z^{qm}(t), \quad t > a. \quad (4.46)$$

Moreover, it is clear that as φ and v are nonnegative continuous functions in $[a, \infty)$.

Thus w is a continuous, nonnegative and nondecreasing function in $[a, \infty)$.

Now, we would like to estimate the right hand side of (4.46) in term of $w(t)$. From (4.44) and (4.45) we entail that

$$z^q(t) \leq 2^{q-1} \left(|b|^q \left(\log \frac{t}{a} \right)^{q\beta(1-\alpha)} + w(t) \right), \quad t > a. \quad (4.47)$$

Raising both sides of (4.47) to the power m and using Lemma 2.5.2 again, we get

$$z^{qm}(t) \leq 2^{mq-1} \left(|b|^{mq} \left(\log \frac{t}{a} \right)^{mq\beta(1-\alpha)} + w^m(t) \right), \quad t > a. \quad (4.48)$$

The substitution of (4.48) in (4.46) yields

$$\begin{aligned} w'(t) &\leq 2^{mq-1} \hat{C}_1^q \varphi^q(t) \left(\log \frac{t}{a} \right)^{-qm\beta(1-\alpha)} \left(|b|^{mq} \left(\log \frac{t}{a} \right)^{mq\beta(1-\alpha)} + w^m(t) \right) \\ &\leq 2^{mq-1} |b|^{mq} \hat{C}_1^q \varphi^q(t) + 2^{mq-1} \hat{C}_1^q \left(\log \frac{t}{a} \right)^{-qm\beta(1-\alpha)} \varphi^q(t) w^m(t). \end{aligned} \quad (4.49)$$

Applying Lemma 2.5.3 (with $w(u) = u^m$) we infer that

$$w(t) \leq G^{-1} \left[G \left(w(a) + \int_a^t 2^{mq-1} |b|^{mq} \hat{C}_1^q \varphi^q(s) ds \right) \right]$$

$$+ \int_a^t 2^{mq-1} \hat{C}_1^q \left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} \varphi^q(s) ds \Big].$$

Let us set

$$l(t) = 2^{mq-1} |b|^{mq} \hat{C}_1^q \int_a^t \varphi^q(s) ds$$

and

$$k(t) = 2^{mq-1} \hat{C}_1^q \int_a^t \left(\log \frac{s}{a} \right)^{-qm\beta(1-\alpha)} \varphi^q(s) ds,$$

then

$$w(t) \leq G^{-1} [G(l(t)) + k(t)],$$

where we have used the fact that $w(a) = 0$. Since $G(r) = \int_{r_0}^r \frac{ds}{s^m}$, $r > 0$, $r_0 > 0$, then

$G(r) = \frac{r^{1-m}}{1-m} - \frac{r_0^{1-m}}{1-m}$ and $G^{-1}(y) = [r_0^{1-m} - (m-1)y]^{-\frac{1}{m-1}}$. That is

$$\begin{aligned} w(t) &\leq G^{-1} \left[\frac{l(t)^{1-m}}{1-m} - \frac{l(t_0)^{1-m}}{1-m} + k(t) \right] \\ &\leq \left[l(t_0)^{1-m} - (m-1) \left(\frac{l(t)^{1-m}}{1-m} - \frac{l(t_0)^{1-m}}{1-m} + k(t) \right) \right]^{-\frac{1}{m-1}} \\ &\leq [l(t)^{1-m} - (m-1)k(t)]^{-\frac{1}{m-1}} \end{aligned}$$

as long as

$$l(t)^{m-1} k(t) < \frac{1}{m-1}.$$

In particular, if $\left(\|\varphi(t)\|_q \right)^{m-1} \left\| \varphi(t) \left(\log \frac{t}{a} \right)^{-m\beta(1-\alpha)} \right\|_q < \mathbf{L}^*/2$, then $w(t) \leq K_1$ for

some positive constant K_1 for all $t > a$, and thus from (4.43) we see that

$$z(t) \leq |b| \left(\log \frac{t}{a} \right)^{\beta(1-\alpha)} + K_1^{1/q},$$

and then

$$v(t) \leq |b| + K_1^{1/q} \left(\log \frac{t}{a} \right)^{-\beta(1-\alpha)} \leq C, \quad t \geq t_0 > ae$$

for some positive constant C . This yields that $|u(t)| \leq C \left(\log \frac{t}{a} \right)^{\gamma-1}$ for $t \geq t_0 > ae$.

The proof is complete.

4.4 Non-existence result

We consider the Cauchy-type problem

$$\begin{cases} \left(\mathcal{D}_{a^+}^{\alpha, \beta} u \right)(t) = f[t, u(t)], & t > a \\ \left(\mathcal{D}_{a^+}^{(\beta-1)(1-\alpha)} u \right)(a) = b \end{cases} \quad (4.50)$$

where

$$\left(\mathcal{D}_{a^+}^{\alpha, \beta} u \right)(t) = \mathcal{J}_{a^+}^{\beta(1-\alpha)} \left(t \frac{d}{dt} \right) \left(\mathcal{J}_{a^+}^{(1-\beta)(1-\alpha)} u \right)(t)$$

is the Hilfer-Hadamard Fractional Derivative (**HHFD**) of order $0 < \alpha < 1$ and type

$$0 \leq \beta \leq 1.$$

We investigate the case when $f[t, u(t)] \geq \left(\log \frac{t}{a} \right)^\mu |u(t)|^m$ for some $m > 1$ and $\mu \in \mathbb{R}$.

That is we consider the Cauchy problem

$$\begin{cases} \left(\mathcal{D}_{a+}^{\alpha,\beta} u \right) (t) \geq \left(\log \frac{t}{a} \right)^\mu |u(t)|^m, & t > a > 0, \ m > 1, \ \mu \in \mathbb{R} \\ \left(\mathcal{D}_{a+}^{\gamma-1} u \right) (a) = b \geq 0 \end{cases} \quad (4.51)$$

where $\mathcal{D}_{a+}^{\alpha,\beta}$ is the **HHFD** of order $0 < \alpha < 1$ and type $0 \leq \beta \leq 1$ and $\gamma = \alpha + \beta - \alpha\beta$.

For this we need the following lemma.

Lemma 4.4.1: *If $\alpha > 0$ and $f \in C[a, b]$, then*

$$\left(\mathcal{J}_{a+}^\alpha f \right) (a) = \lim_{t \rightarrow a} \left(\mathcal{J}_{a+}^\alpha f \right) (t) = 0$$

and

$$\left(\mathcal{J}_{b-}^\alpha f \right) (b) = \lim_{t \rightarrow b} \left(\mathcal{J}_{b-}^\alpha f \right) (t) = 0.$$

Proof : Since $f \in C[a, b]$, then on $[a, b]$, we have

$$|f(t)| < M,$$

for some positive constant M .

Therefore

$$\begin{aligned} \left| \left(\mathcal{J}_{a+}^\alpha f \right) (t) \right| &\leq \frac{1}{\Gamma(\alpha)} \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} |f(s)| \frac{ds}{s} \\ &\leq \frac{M}{\Gamma(\alpha)} \int_a^t \left(\log \frac{t}{s} \right)^{\alpha-1} \frac{ds}{s} \\ &\leq \frac{M}{\alpha \Gamma(\alpha)} \left[- \left(\log \frac{t}{s} \right)^\alpha \right]_{s=a}^t = \frac{M}{\Gamma(\alpha+1)} \left(\log \frac{t}{a} \right)^\alpha. \end{aligned}$$

As $\alpha > 0$ we see that

$$(\mathcal{J}_{a+}^\alpha f)(a) = \lim_{t \rightarrow a} (\mathcal{J}_{a+}^\alpha f)(t) = 0.$$

Similarity

$$\begin{aligned} |(\mathcal{J}_{b-}^\alpha f)(t)| &\leq \frac{1}{\Gamma(\alpha)} \int_t^b \left(\log \frac{s}{t}\right)^{\alpha-1} |f(s)| \frac{ds}{s} \\ &\leq \frac{M}{\Gamma(\alpha)} \int_t^b \left(\log \frac{s}{t}\right)^{\alpha-1} \frac{ds}{s} \\ &\leq \frac{M}{\alpha \Gamma(\alpha)} \left[\left(\log \frac{s}{t}\right)^\alpha \right]_{s=t}^b = \frac{M}{\Gamma(\alpha+1)} \left(\log \frac{b}{t}\right)^\alpha. \end{aligned}$$

As $\alpha > 0$ we see that

$$(\mathcal{J}_{b-}^\alpha f)(b) = \lim_{t \rightarrow b} (\mathcal{J}_{b-}^\alpha f)(t) = 0.$$

Theorem 4.4.1: Assume that $\mu \geq 0$. Then, Problem (4.51) does not admit global nontrivial solution in $C_{1-\gamma, \log}^\gamma[a, b]$ when $b \geq 0$.

Proof : Assume that a nontrivial solution exists for all time $t > a$. Let $\varphi(t) \in C^1([a, \infty))$ be a test function satisfying : $\varphi(t) \geq 0$, $\varphi(t)$ is non-increasing and such that

$$\varphi(t) := \begin{cases} 1, & a \leq t \leq \theta T \\ 0, & t \geq T \end{cases}$$

for some $T > a$ and some θ ($\theta < 1$) such that $a < \theta T < T$. Multiplying the inequality in (4.51) by $\frac{\varphi(t)}{t}$ and integrating over $[a, T]$ we get

$$\int_a^T \varphi(t) \left(\mathcal{D}_{a+}^{\alpha, \beta} u \right)(t) \frac{dt}{t} \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}$$

and from the definition of $\left(\mathcal{D}_{a+}^{\alpha,\beta}u\right)(t)$ we can write

$$\int_a^T \varphi(t) \left(\mathcal{J}_{a+}^{\beta(1-\alpha)} t \frac{d}{dt} \mathcal{J}_{a+}^{1-\gamma} u \right)(t) \frac{dt}{t} \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \quad (4.52)$$

By virtue of Lemma 2.3.6 (after extending by zero outside $[a, T]$), we may deduce from (4.52) that

$$\int_a^T \frac{d}{dt} \left(\mathcal{J}_{a+}^{1-\gamma} u \right)(t) \left(\mathcal{J}_{T-}^{\beta(1-\alpha)} \varphi(t) \right)(t) dt \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \quad (4.53)$$

Notice that Lemma 2.3.6 is valid in our case since $\left(\left(\log \frac{t}{a} \right)^{(1-\gamma)} (\mathcal{D}_{a+}^\gamma u) \right) \in C[a, T]$ implies that $\left| \left(\log \frac{t}{a} \right)^{(1-\gamma)} (\mathcal{D}_{a+}^\gamma u)(t) \right| \leq M$ on $[a, T]$ for some positive constant M

$$\begin{aligned} \int_a^T \left| t^{-\frac{1}{p}} (\mathcal{D}_{a+}^\gamma u)(t) \right|^{p'} \frac{dt}{t} &\leq M \int_a^T t^{1-p'} \left(\log \frac{t}{a} \right)^{-p'(1-\gamma)} \frac{dt}{t} \\ &\leq M \int_a^\infty t^{1-p'} \left(\log \frac{t}{a} \right)^{-p'(1-\gamma)} \frac{dt}{t}. \end{aligned}$$

Let $s = (p' - 1) \left(\log \frac{t}{a} \right)$, then by the definition of the Gamma function

$$\begin{aligned} \int_a^T \left| t^{-\frac{1}{p}} (\mathcal{D}_{a+}^\gamma u)(t) \right|^{p'} \frac{dt}{t} &\leq \frac{M a^{1-p'}}{(p' - 1)^{1-p'(1-\gamma)}} \int_0^\infty s^{-p'(1-\gamma)} e^{-s} ds \\ &\leq \frac{M a^{1-p'}}{(p' - 1)^{1-p'(1-\gamma)}} \Gamma(1 - p'(1 - \gamma)) < \infty. \end{aligned}$$

Hence $t \frac{d}{dt} (\mathcal{J}_{a^+}^{1-\gamma} u) (t) = (\mathcal{D}_{a^+}^\gamma u) (t) \in X_{-1/p}^{p'}$ (and $\varphi \in L_p$) for some $p > \frac{1}{\gamma}$.

An integration by parts in (4.53) yields

$$\begin{aligned} & \left[(\mathcal{J}_{a^+}^{1-\gamma} u) (t) (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (t) \right]_{t=a}^T - \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u) (t) \frac{d}{dt} (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (t) dt \\ & \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t} \end{aligned}$$

or

$$\begin{aligned} & -b (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (a^+) - \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u) (t) \frac{d}{dt} (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (t) dt \\ & \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t} \end{aligned} \quad (4.54)$$

because $(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (T) = 0$ (see Lemma 4.4.1) and

$$(\mathcal{J}_{a^+}^{1-\gamma} u) (a) = (\mathcal{D}_{a^+}^{\gamma-1} u) (a) = b.$$

Multiplying by $\frac{t}{t}$ inside the integral in the left hand side of (4.54) we see that

$$\begin{aligned} & -b (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (a) + \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u) (t) \left(-t \frac{d}{dt} \right) (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (t) \frac{dt}{t} \\ & \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \end{aligned}$$

It appears from Definition 2.3.4 that

$$-b (\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi) (a^+) + \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u) (t) (\mathcal{D}_{T^-}^{1-\beta(1-\alpha)} \varphi) (t) \frac{dt}{t}$$

$$\geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}$$

and from Lemma 2.3.3 we see that

$$\begin{aligned} & -b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) + \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u)(t) \left[\frac{\varphi(T)}{\Gamma(\beta(1-\alpha))} \left(\log \frac{T}{t} \right)^{\beta(1-\alpha)-1} \right. \\ & \left. - \frac{1}{\Gamma(\beta(1-\alpha))} \int_t^T \left(\log \frac{s}{t} \right)^{\beta(1-\alpha)-1} \varphi'(s) ds \right] \frac{dt}{t} \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \end{aligned}$$

Since $\varphi(T) = 0$ and

$$\frac{1}{\Gamma(\beta(1-\alpha))} \int_t^T \left(\log \frac{s}{t} \right)^{\beta(1-\alpha)-1} \varphi'(s) ds = \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \delta \varphi \right) (t),$$

the last inequality becomes

$$\begin{aligned} & -b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) - \int_a^T (\mathcal{J}_{a^+}^{1-\gamma} u)(t) \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \delta \varphi \right) (t) \frac{dt}{t} \\ & \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \end{aligned}$$

Note that $\delta \varphi \in L_p$ and by the same argument as the one used at the beginning of the proof we may show that $\mathcal{J}_{a^+}^{1-\gamma} u \in X_{-1/p}^{p'}$ since $\mathcal{J}_{a^+}^{1-\gamma} u \in C_{1-\gamma, \log} [a, T]$. Therefore,

Lemma 2.3.6 again allows us to write

$$\begin{aligned} & -b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) - \int_a^T \delta \varphi(t) \left(\mathcal{J}_{a^+}^{\beta(1-\alpha)} \mathcal{J}_{a^+}^{1-\gamma} u \right) (t) \frac{dt}{t} \\ & \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}, \end{aligned}$$

and by the Semigroup Property 3.3.2

$$-b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) - \int_a^T \delta \varphi(t) \left(\mathcal{J}_{a^+}^{1-\alpha} u \right) (t) \frac{dt}{t} \geq \int_a^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t}. \quad (4.55)$$

On the other hand

$$\begin{aligned} \int_a^T \delta \varphi(t) \left(\mathcal{J}_{a^+}^{1-\alpha} u \right) (t) \frac{dt}{t} &= \frac{1}{\Gamma(1-\alpha)} \int_a^T \delta \varphi(t) \int_a^t \left(\log \frac{t}{s} \right)^{-\alpha} \frac{u(s)}{s} ds \frac{dt}{t} \\ &\leq \frac{1}{\Gamma(1-\alpha)} \int_a^T |\delta \varphi(t)| \int_a^t \left(\log \frac{t}{s} \right)^{-\alpha} \frac{|u(s)|}{s} ds \frac{dt}{t}. \end{aligned}$$

As φ is nonincreasing, we have $\varphi(s) \geq \varphi(t)$ for all $t \geq s$ and $\frac{1}{\varphi^{1/m}(s)} \leq \frac{1}{\varphi^{1/m}(t)}$, $m > 1$.

Also it is clear that

$$\varphi'(t) = 0, \quad t \in [a, \theta T].$$

Therefore

$$\begin{aligned} &\int_a^T \delta \varphi(t) \left(\mathcal{J}_{a^+}^{1-\alpha} u \right) (t) \frac{dt}{t} \\ &\leq \frac{1}{\Gamma(1-\alpha)} \int_a^T |\delta \varphi(t)| \int_a^t \left(\log \frac{t}{s} \right)^{-\alpha} \frac{|u(s)| \varphi^{1/m}(s)}{s \varphi^{1/m}(s)} ds \frac{dt}{t} \\ &\leq \frac{1}{\Gamma(1-\alpha)} \int_{\theta T}^T \frac{|\delta \varphi(t)|}{\varphi^{1/m}(t)} \int_a^t \left(\log \frac{t}{s} \right)^{-\alpha} \frac{|u(s)| \varphi^{1/m}(s)}{s} ds \frac{dt}{t}. \end{aligned}$$

Definition 2.3.1 allows us to write

$$\int_a^T \delta \varphi(t) \left(\mathcal{J}_{a^+}^{1-\alpha} u \right) (t) \frac{dt}{t} \leq \int_{\theta T}^T \frac{|\delta \varphi(t)|}{\varphi^{1/m}(t)} \left(\mathcal{J}_{a^+}^{1-\alpha} |u| \varphi^{1/m} \right) (t) \frac{dt}{t}.$$

By the same argument as the one used at the beginning of the proof we may show that $|u(t)| \varphi^{1/m}(t) \in X_{-1/p}^{p'} (|u(t)| \varphi^{1/m}(t) < |u(t)|)$. Moreover, it is easy to see that $\frac{|\delta\varphi(t)|}{\varphi^{1/m}(t)} \in L_p$ (for otherwise we consider $\varphi^\lambda(t)$ with some sufficiently large λ). Thus, we can apply Lemma 2.3.6 to get

$$\int_a^T \delta\varphi(t) (\mathcal{J}_{a^+}^{1-\alpha} u)(t) \frac{dt}{t} \leq \int_{\theta T}^T |u(t)| \varphi^{1/m}(t) \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta\varphi|}{\varphi^{1/m}} \right)(t) \frac{dt}{t}. \quad (4.56)$$

Next, we multiply by $(\log \frac{t}{a})^{\mu/m} \cdot (\log \frac{t}{a})^{-\mu/m}$ inside the integral in the right hand side of (4.56)

$$\int_a^T \delta\varphi(t) (\mathcal{J}_{a^+}^{1-\alpha} u)(t) \frac{dt}{t} \leq \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta\varphi|}{\varphi^{1/m}} \right)(t) |u(t)| \varphi^{1/m}(t) \frac{(\log \frac{t}{a})^{\mu/m}}{(\log \frac{t}{a})^{\mu/m}} \frac{dt}{t}.$$

For $\mu \geq 0$ we have $(\log \frac{t}{a})^{-\mu/m} \leq (\log \frac{\theta T}{a})^{-\mu/m}$ (because $-\mu/m < 0$ and $t > \theta T$), that is

$$\left(\log \frac{t}{a} \right)^{-\mu/m} \leq \left(\log \frac{\theta T}{a} \right)^{-\mu/m}.$$

It follows that

$$\begin{aligned} & \int_a^T \delta\varphi(t) (\mathcal{J}_{a^+}^{1-\alpha} u)(t) \frac{dt}{t} \\ & \leq \left(\log \frac{\theta T}{a} \right)^{-\mu/m} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta\varphi|}{\varphi^{1/m}} \right)(t) \left(\log \frac{t}{a} \right)^{\mu/m} |u(t)| \varphi^{1/m}(t) \frac{dt}{t}. \end{aligned} \quad (4.57)$$

By using the Young inequality (see Theorem 2.5.2), with m and m' such that $\frac{1}{m} + \frac{1}{m'} = 1$, in the right hand side of (4.57) we find

$$\int_a^T \delta\varphi(t) (\mathcal{J}_{a^+}^{1-\alpha} u)(t) \frac{dt}{t}$$

$$\begin{aligned}
&\leq \frac{1}{m} \int_{\theta T}^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} + \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{m'} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right)^{m'}(t) \frac{dt}{t} \\
&\leq \frac{1}{m} \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} + \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{m'} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right)^{m'}(t) \frac{dt}{t}.
\end{aligned} \tag{4.58}$$

Clearly from (4.55) and (4.58) we see that

$$\begin{aligned}
&-b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) + \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{m'} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right)^{m'}(t) \frac{dt}{t} \\
&\geq \left(1 - \frac{1}{m} \right) \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t},
\end{aligned}$$

or, since $b \geq 0$ and $\varphi(t) \geq 0$ ($b \left(\mathcal{J}_{T^-}^{\beta(1-\alpha)} \varphi \right) (a^+) \geq 0$)

$$\frac{1}{m'} \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \leq \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{m'} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right)^{m'}(t) \frac{dt}{t}.$$

Therefore, by Definition 2.3.2 we have

$$\begin{aligned}
&\int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \\
&\leq \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{\Gamma^{m'}(1-\alpha)} \int_{\theta T}^T \left(\int_t^T \left(\log \frac{s}{t} \right)^{-\alpha} \frac{|\delta \varphi(s)|}{\varphi^{1/m}(s)} \frac{ds}{s} \right)^{m'} \frac{dt}{t}.
\end{aligned}$$

The change of variable $\sigma T = t$ yields

$$\int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \leq \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{\Gamma^{m'}(1-\alpha)} \int_\theta^1 \left(\int_{\sigma T}^T \left(\log \frac{s}{\sigma T} \right)^{-\alpha} \frac{|\varphi'(s)|}{\varphi(s)^{1/m}} ds \right)^{m'} \frac{d\sigma}{\sigma}.$$

Another change of variable $r = s/T$ gives

$$\begin{aligned} & \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \\ & \leq \frac{\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m}}{\Gamma^{m'}(1-\alpha)} \int_\theta^1 \left(\int_\sigma^1 \left(\log \frac{r}{\sigma} \right)^{-\alpha} \frac{|\varphi'(r)|}{\varphi(r)^{1/m}} dr \right)^{m'} \frac{d\sigma}{\sigma}. \end{aligned} \quad (4.59)$$

We may assume that the integral term in the right hand side of (4.59) is convergent, that is

$$\frac{1}{\Gamma^{m'}(1-\alpha)} \int_\theta^1 \left(\int_\sigma^1 \left(\ln \frac{r}{\sigma} \right)^{-\alpha} \frac{|\varphi'(r)|}{\varphi(r)^{1/m}} dr \right)^{m'} d\sigma \leq C$$

for some positive constant C , for otherwise we consider $\varphi^\lambda(r)$ with some sufficiently large λ . Therefore

$$\int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \leq C \left(\log \frac{\theta T}{a} \right)^{-\mu m'/m} \quad (4.60)$$

If $\mu > 0$, we claim that $-\frac{\mu m'}{m} < 0$, and consequently

$$\left(\log \frac{\theta T}{a} \right)^{-\mu m'/m} \rightarrow 0$$

as $T \rightarrow \infty$. Finally from (4.60) we obtain

$$\lim_{T \rightarrow \infty} \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} = 0.$$

We reach a contradiction since the solution is not supposed to be trivial.

In the case $\mu = 0$ we have $-\mu m'/m = 0$ and the relation (4.60) ensures that

$$\lim_{T \rightarrow \infty} \int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \leq C. \quad (4.61)$$

Moreover, it is clear that

$$\begin{aligned} & \left(\log \frac{\theta T}{a} \right)^{-\mu/m} \int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right) (t) \left(\log \frac{t}{a} \right)^{\mu/m} |u(t)| \varphi^{1/m}(t) \frac{dt}{t} \\ & \leq \left(\log \frac{\theta T}{a} \right)^{-\mu/m} \left[\int_{\theta T}^T \left(\mathcal{J}_{T^-}^{1-\alpha} \frac{|\delta \varphi|}{\varphi^{1/m}} \right)^{m'} (t) \frac{dt}{t} \right]^{\frac{1}{m'}} \left[\int_{\theta T}^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t} \right]^{\frac{1}{m}}. \end{aligned}$$

This relation, together with (4.55) and (4.57), implies that

$$\int_a^T \left(\log \frac{t}{a} \right)^\mu \varphi(t) |u(t)|^m \frac{dt}{t} \leq K \left[\int_{\theta T}^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t} \right]^{\frac{1}{m}}$$

for some positive constant K , with

$$\lim_{T \rightarrow \infty} \int_{\theta T}^T \left(\log \frac{t}{a} \right)^\mu |u(t)|^m \varphi(t) \frac{dt}{t} = 0$$

due to the convergence of the integral in (4.61). This is again a contradiction. This completes the proof of Theorem 3.4.1.

Chapter 5

RECOMMENDATIONS

In this thesis we have restricted ourselves to the case $0 \leq \alpha, \beta \leq 1$. It will be certainly interesting to extend the Hilfer fractional derivative and Hilfer-Hadamard fractional derivative to the cases where α and β takes values above one.

It is also important to consider different types of nonlinearities in the stability and non-existence issues other than the ones we have investigated in this document.

Due to the time constraint the Hilfer fractional derivative has been defined and studied only for the Hadamard fractional derivative. It is worth introducing a similar Hilfer type fractional derivative which interpolates the Erdélyi-Kober fractional derivative and the corresponding Caputo type modification of the Erdélyi-Kober fractional derivative.

Bibliography

- [1] S. A. Abd El-Salam, On the stability of some fractional-order non-autonomous systems, *Electronic Journal of Qualitative Theory of Differential Equations*. No. 6, (2007), 1-14.
- [2] B. Ahmad and S. Sivasundaram, Some existence results for fractional integro-differential equations with nonlinear conditions, *Communications in Applied Analysis* 12, no. 2, (2008), 107-112.
- [3] B. Ahmad and S. Sivasundaram, Some basic results for fractional functional integro-differential equations. *Communications in Applied Analysis* 12, no. 4, (2008), 467-478.
- [4] A. Z. Al-Abedeen, Existence theorem on differential equation of generalized order, *Rafidain J. Sci. Mosul. Univ. Iraq*, 1, (1976), 95-104.
- [5] A. Z. Al-Abedeen and H. L. Arora, A global existence and uniqueness theorem for ordinary differential equation of generalized order, *Canad. Math. Bull.*, 21, n. 3, (1978), 267-271.

- [6] M. A. Al-Bassam, Some existence theorems on differential equations of generalized order, *J. Reine Angew. Math.*, 218, (1965), 70-78.
- [7] H. L. Arora and J. G. Alshamari, Stability of differential equations of non-integer order through fixed point in the large, *Indian J. Pure Appl. Math.*, 11, n. 3, (1980), 307-313.
- [8] Yu. Babenko, Heat and Mass Transfer, *Chimia*, Leningrad 1986. [in Russian].
- [9] R. L. Bagley and P. J. Torvik, A theoretical basis for the application of fractional calculus to viscoelasticity, *J. Rheology* 27, (1983), 201-210.
- [10] R. L. Bagley and P. J. Torvik, A different approach to the analysis of viscoelastically damped structures, *AIAA Journal* 21, (1983), 741-748.
- [11] R. L. Bagley and P. J. Torvik, On the appearance of the fractional derivative in the behavior of real material, *J. Appl. Mechanics* 51, (1983), 294-298.
- [12] R. L. Bagley and R. A. Calico, Fractional order state equations for the control of viscoelastic structures, *Journal of Guidance, Control, and Dynamics* 14(2), 1991, 304-311.
- [13] M. Benchohra, S. Hamani and S. K. Ntouyas, Boundary value problems for differential equations with fractional order, *Surveys in Mathematics and its Applications*, 3, (2008), 1-12.

- [14] M. Benchohra, J. Henderson, and D. Seba, Measure of noncompactness and fractional differential equations in Banach spaces, *Communications in Applied Analysis* 12, no. 4, (2008), 419-428.
- [15] P. L. Butzer, A. A. Kilbas and J. J. Trujillo, Mellin transform analysis and integration by parts for Hadamard-type fractional integral, *J. Math. Anal. Appl.* 270, no. 1, (2002) 1-15.
- [16] M. Caputo, *Elasticità e Dissipazione*, Zanichelli, Bologna 1969. [in Italian].
- [17] H. T. Davis, *The Theory of Linear Operators*, The Principia Press, Bloomington, Indiana 1936.
- [18] D. Delbosco and L. Rodino, Existence and uniqueness for a nonlinear fractional differential equation, *J. Math. Anal. Appl.*, 204, no. 2, (1996), 609-625.
- [19] J. V. Devi, F. A. McRae, Z. Drici, Generalized quasilinearization for fractional differential equations, *Computers and Mathematics with Applications*, 59, (2010), 1057-1062.
- [20] J. V. Devi and CH. Suseela, Quasilinearization for fractional differential equations, *Communications in Applied Analysis* 12, no. 4, (2008), 407-418.
- [21] K. Diethelm and A. D. Freed, On the solution of nonlinear fractional order differential equations used in the modeling of viscoplasticity, in "Science Computing in Chemical Engineering II-Computational Fluid Dynamics, Reaction Engineering and Molecular Properties" (F. Keil, W. Mackens, H. Voss, and J. Werther, Eds), pp 217-224, Springer-Verlag, Heidelberg, 1999.

- [22] K. Diethelm and N. J. Ford, Analysis of fractional differential equations, J. Math. Anal. Appl. 265 (2002), 229-248.
- [23] K. Diethelm and G. Walz, Numerical solution of fractional order differential equations by extrapolation, Numer. Algorithms 16 (1997), 231-253.
- [24] K. Diethelm and N. J. Ford, Analysis of Fractional Differential Equations, Manchester Centre for Computational Mathematics Numerical Analysis Reports, (2003).
- [25] M. M. Dzherbashian, Integral Transforms and Representations of Functions in the Complex Plane, Nauka, Moscow 1966. [in Russian].
- [26] M. M. Dzherbashian, Harmonic Analysis and Boundary Value Problems in the Complex Domain, Birkhauser Verlag, Basel 1993.
- [27] A. M. A. El-Sayed. Fractional differential equations, Kyungpook Math. J., 28, n. 2, (1988), 119-122.
- [28] A. M. A. El-Sayed, On the fractional differential equations, Appl. Math. Comput., 49, n. 2-3, (1992), 205-213.
- [29] A. M. A. El-Sayed, Linear differential equations of fractional order, Appl. Math. Comput., 55, n. 1, (1993), 1-12.
- [30] A. M. A. El-Sayed, Finite Weyl fractional calculus and abstract fractional differential equation, J. Fract. Calc., 9, (1996), 59-68.

- [31] A. M. A. El-Sayed and F. M. Gafar, Exact solution of linear differential equations of fractional order, *Kyungpook Math. J.*, 37, n. 2. (1997), 181-189.
- [32] A. M. A. El-Sayed and F. M. Gafar, Sturm-Liouville equation of fractional order, *Kyungpook Math. J.*, 37, n. 1, (1997), 9-17.
- [33] A. M. A. El-Sayed and A.-G. Ibragim, Multivalued fractional differential equations, *Appl. Math. Comput.*, 68, n. 1, (1995), 15-25.
- [34] A. M. A. El-Sayed, Sh. A. Abd El-Salam, Weighted Cauchy-type problem of a functional differ-integral equation, *Electronic Journal of Qualitative Theory of Differential Equations*, no. 30, (2007), 1-9.
- [35] A. M. A. El-Sayed, Sh. A. Abd El-Salam, solution of Weighted Cauchy-type Problem of a Diffre-integral Functional Equation, *International Journal of Non-linear Science*, Vol. 5, no. 3, (2008), 281-288.
- [36] A. Erdélyi, *Tables of Integral Transforms*, Bateman Project, Vols. 1-2, McGraw-Hill, New York 1953-1954.
- [37] K. M. Furati and N-e. Tatar, Power type estimates for a nonlinear fractional differential equation, *Nonlinear Analysis T.M.A.* 62 (2005), 1025-1036.
- [38] K. M. Furati and N-e. Tatar, An existence result for a nonlocal fractional differential problem, *J. Fract. Calc.* Vol. 26 (2004), 43-51.
- [39] K. M. Furati and N-e. Tatar, Behavior of solutions for a weighted Cauchy-type fractional differential problem, *J. Fract. Calc.* 28 (2005), 23-42.

- [40] K. M. Furati and N-e. Tatar, Long time behaviour for a nonlinear fractional model, *Journal of Mathematical Analysis and Applications*, Vol. 332, Issue 1 (2007), 441-454.
- [41] K. M. Furati and N- e. Tatar, Some fractional differential inequalities and their applications, *Mathematical Inequalities and Applications*, Vol. 9, Issue 4 (2006), 577-598.
- [42] L. Gaul, P. Klein and S. Kempfle, Damping description involving fractional operators, *Mech. Systems Signal Processing* 5 (1991), 81-88.
- [43] I. M. Gelfand and G.E. Shilov, *Generalized Functions*, Vol. 1, Academic Press, New York, 1964.
- [44] W. G. Glöckle and T. F. Nonnenmacher, A fractional calculus approach of selfsimilar protein dynamics, *Biophys. J.* 68 (1995), 46-53.
- [45] R. Gorenflo and S. Vessella, *Abel Integral Equations: Analysis and Applications*, *Lecture Notes in Mathematics*, Springer-Verlag, Berlin, 1991.
- [46] T. T. Hartley, C. F. Lorenzo and H. K. Qammar, Chaos in a fractional order Chua system, *IEEE Transactions on Circuits & Systems: Part I* 42(8), 1995, 485-490.
- [47] N. Hayek, J. Trujillo, M. Rivero, B. Bonilla and J. C. Moreno, An extension of Picard-Lindeloff theorem to fractional differential equations, *Appl. Anal.*, 70, n. 3-4, (1999), 347-361.

- [48] O. Heaviside, Electromagnetic Theory, Vol. II, Chelsea Edition (1971), New York, 1922.
- [49] R. Hilfer, Fractional time evolution, Applications of fractional calculus in Physics, World Scientific, New-Jersey, London-Hong Kong, 2000, 87-130.
- [50] M. Ichise, Y. Nagayanagi and T. Kojima, An analog simulation of non-integer order transfer functions for analysis of electrode processes, J. Electroanalytical Chemistry Interfacial Electrochemistry 33, 1971, 253-265.
- [51] A. A. Kilbas, B. Bonilla and J.J. Trujillo, Existence and uniqueness theorems for nonlinear fractional differential equations, Demonstratio Math., 33, n. 3, (2000), 583-602.
- [52] A. A. Kilbas, B. Bonilla and J. J. Trujillo, Fractional integrals and derivatives and differential equations of fractional order in weighted spaces of continuous functions. (Russian), Dokl. Nats. Akad. Nauk Belarusi, 44, n. 6, (2000).
- [53] A. A. Kilbas and S. A. Marzan, Cauchy problem for differential equation with Caputo derivative, An International Journal for Theory and applications, 7, n. 3, (2004), 297-320.
- [54] A. A. Kilbas and S. A. Marzan, Nonlinear differential equations with the Caputo fractional derivative in the space of continuously differentiable functions, Differential equations, Vol. 41, No. 1, (2005), 84-89.

- [55] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies 204, Editor: Jan van Mill, Elsevier, Amsterdam, The Netherlands 2006.
- [56] A. A. Kilbas and A. A. Titoura, Nonlinear differential equations with Marchaud-Hadamard-type fractional derivative in the weighted space of summable functions, Mathematical Modelling and Analysis, 12, n. 3, (2007), 243-256.
- [57] R. C. Koeller, Application of fractional calculus to the theory of viscoelasticity, J. Appl. Mechanics 51, (1984), 299-307.
- [58] R. C. Koeller, Polynomial operators, Stieltjes convolution, and fractional calculus in hereditary mechanics, Acta Mechanica 58, (1986), 251-264.
- [59] N. Kosmatov, Integral equations and initial value problems for nonlinear differential equations of fractional order, Nonlinear Analysis, 70, (2009), 2521-2529.
- [60] C. Kou, J. Liu and Y. Ye, Existence and uniqueness of solutions for the Cauchy-type problems of fractional differential equations, Discrete Dynamics in Nature and Society, Volume 2010, Article ID 142175, 2010, 1-15.
- [61] V. Lakshmikantham, S. Leela, Nagumo-type uniqueness result for fractional differential equations, Nonlinear Analysis, 71, (2009), 2886-2889.
- [62] V. Lakshmikantham, S. Leela, J. Vasundhara Devi, Theory of Fractional Dynamic Systems, Cambridge Scientific Publishers, 2009.

- [63] Y. Laskri and N. M. Tatar, "The critical exponent for an ordinary fractional differential problem," *Computers & Mathematics with Applications*, 59, (2010), 1266-1270, .
- [64] I. P. Leskovskii, On the theory of fractional differentiation. (Russian), *Differentsial'nye Uravnenija*, 13, n. 1, (1977), 170-173.
- [65] Wei Lin, Global existence theory and chaos control of fractional differential equations, *Journal of Mathematical Analysis and applications*, 332, (2007), 709-726.
- [66] Z. Luszczki and B. Rzepecki, An existence theorem for ordinary differential equations of order $\alpha \in (0, 1]$, *Demonstratio Math.*, 20, n. 3-4, (1987), 471-475 (1988).
- [67] F. Mainardi, Fractional calculus: Some basic problems in continuum and statistical mechanis, in "Fractals and Fractional Calculus in Continuum Mechanics" (A. Carpinteri and F. Mainardi, Eds), pp. 291-348, Springer-Verlag, Wien, 1997.
- [68] F. Mainardi, Fractional Calculus and Waves in Linear Viscoelastsity, an Introduction to Mathematical Model, Imperial College Press. World Scientific Publishing, London 2010.
- [69] B. Mandelbrot, Some noises with $1/f$ spectrum, a bridge between direct current and white noise, *IEEE Transactions on Information Theory* 13(2), (1967), 289-298.

- [70] S. Mandelbrojt, Sulla genemlizzazione del calcolo delle variazione. (French),
Atti Reale Accad. Naz. Lincei. Rend. C1. Sci. Fiz. Mat. S. Natur., Ser. 6,1,
(1925), 151-156.
- [71] F. Metzler, W. Schick, H. G. Kilian and T. F. Nonnenmacher, Relaxation in
filled polymers: A fractional calculus approach, J. Chem. Phys. 103 (1995),
7180-7186.
- [72] M. W. Michalski, Derivatives of non integer order and their applications, Dis-
sertationes Mathematicae, Polska Akademia Nauk, Instytut Matematyczny-
Warszawa, 1993.
- [73] K. S. Miller and B. Ross, An Introduction to the Fractional Calculus and Dif-
ferential Equations, John Wiley, New York, 1993.
- [74] E. Mitidieri, S.I. Pohozaev, A priori estimates and blow-up of solutions to non-
linear partial differential equations and inequalities. Proc. Steklov Inst. Math.,
234, (2001), 1-383.
- [75] S. M. Momani and S. B. Hadid, Some comparison results for integro-fractional
differential inequalities. J. Fract. Calc. 24 (2003), 37-44.
- [76] S. M. Momani, S. B. Hadid and Z. M. Alawenh, Some analytical properties of
solutions of differential equations of noninteger order, Int. J. Math. Math. Sci.
2004 (2004), 697-701.
- [77] S. Momani and S. Hadid, Lyapunov stability solutions of fractional integrodif-
ferential equations, Int. J. Math. Math. Sc., 47, (2004), 2503-2507.

- [78] G. M. N'Guer'ekata, A Cauchy problem for some fractional abstract differential equation with non local conditions, *Nonlinear Analysis*, 70, (2009), 1873-1876.
- [79] K. B. Oldham and J. Spanier, "The Fractional Calculus", Academic Press, New York, London, 1974.
- [80] L. O'Shaughnessy, Problem # 433, *Amer. Math. Month.*, 25, (1918), 172-173.
- [81] B. G. Pachpatte, *Inequalities for Differential and Integral Equations*, volume 197 of *Mathematics in Science and Engineering*, Acad. Press, San Diego-London, 1998. Edited by W. F. Ames.
- [82] E. Pitcher and W. E. Sewell, Existence theorem for solutions of differential equations of non-integral order, *Bull. Amer. Math. Soc.*, 44, n. 2, (1938), 100-107 (A correction in n. 12, p.888).
- [83] I. Podlubny, *Fractional Differential Equations*, *Mathematics in Sciences and Engineering*. 198, Academic Press, San-Diego, 1999.
- [84] I. Podlubny, I. Petraš, B. M. Vinagre, P. O'Leary and L. Dorčák, Analogue realizations of fractional-order controllers. Fractional order calculus and its applications, *Nonlinear Dynam.* 29 (2002), 281-296.
- [85] I. Podlubny, Geometric and physical interpretation of fractional integration and fractional differentiation, *Fract. Calcul. Anal. Appl.* 5 (4) (2002), 367-386.
- [86] E.L. Post, Discussion of the solution of $(d/dx)^{1/2}y = y/x$ (# problem 433), *Amer. Math. Month.*, 26, (1919), 37-39.

- [87] B. Ross, Fractional Calculus and its Applications, Lecture Notes in Mathematics, Springer Verlag, Berlin 1975.
- [88] B. Rubin, Fractional Integrals and Potentials, Pitman Monographs and Surveys in Pure and Applied Mathematics, Addison Wesley Longman, Harlow 1996.
- [89] S. G. Samko, A. A. Kilbas and O. I. Marichev, "Fractional Integrals and Derivatives: Theory and Applications", Gordon and Breach, 1987. (Trans. from Russian 1993).
- [90] N.P. Semenchuk, A class of differential equations of noninteger order. (Russian), *Differentsial'nye Uravnenija*, 18, n. 10, (1982), 1831-1833.
- [91] Z. Shuqin, Monotone iterative method for initial value problem involving Riemann-Liouville fractional derivatives, *Nonlinear Analysis*, 71, (2009), 2087-2093.
- [92] S. B. Skaar, A. N. Michel and R. K. Miller, Stability of viscoelastic control systems, *IEEE Transactions on Automatic Control* 33(4), (1988), 348-357.
- [93] H. H. Sun, A. A. Abdelwahab and B. Onaral, Linear approximation of transfer function with a pole of fractional order, *IEEE Transactions on Automatic Control* 29(5), (1984), 441-444.
- [94] H. H. Sun, B. Onaral and Y. Tsao, Application of positive reality principle to metal electrode linear polarization phenomena, *IEEE Transactions on Biomedical Engineering* 31(10), 1984, 664-674.

- [95] A. Z.-A. M. Tazali, Local existence theorems for ordinary differential equations of fractional order, Ordinary and Partial Differential Equations (Dundee, 1982), Lecture Notes in Math., 964, (1982), 652-665.
- [96] A. Z. Tazali and I.F. Karim, Continuous dependence on parameter and continuation of solution in the extended sense in ordinary differential equations of fractional order, Dirasat Ser. B. Pure Apl. Sc1, 21, n, 1, (1994), 270-280.
- [97] H. Ye, J. Gao, Y. Ding, A generalized Gronwall inequality and its application to a fractional differential equation, Journal of Mathematical Analysis and applications, 328, (2007), 1075-1081.
- [98] C. Yu., G. Gao, On the solution of nonlinear fractional order differential equation, Nonlinear Analysis, 63, (2005), 971-976.
- [99] C. Yu and G. Gao, Existence of fractional differential equations, Journal of Mathematical Analysis and applications, 310, (2005), 26-29.
- [100] Y. Zhou, Existence and uniqueness of solutions for a system of fractional differential equations, An International Journal for Theory and applications, 12, n.2, (2009), 195-204.

Vita

PERSONAL DATA

Name: Mohammed Dahan Ahmed Kassim

Date of Birth: May. 09th, 1978

Marital Status: Married

Nationality: Yemeni

Institution: King Fahd University of Petroleum and Minerals

Department: Department of Mathematics and Statistics

E-mail: g200705070@kfupm.edu.sa

dahan1978@yahoo.com

HIGHER INSTITUTION ATTENDED

INSTITUTION

1. King Fahd University of Petroleum and Minerals 2007 to date
2. Sana'a university 1999-2002

QUALIFICATIONS

- 1- Master of Science In view
- 2- Bachelor in Mathematics 2002

PROFESSIONAL EXPERIENCE

Teaching as a Lecturer in Sana'a university from 2002 to 2007

COURSES TAUGHT

CALCULUS I, CALCULUS II, SET THEORY, PROBABILITY, PASCAL